

Probability theory

PETER PFAFFELHUBER

Version: April 11, 2024

Prelude

These are the notes of a lecture which I gave at the University of Freiburg. After some elementary probability and measure theory, this course introduces some main concepts in (measure theoretic) probability theory. As a prerequisite, for measure theoretic contents, we refer to my manuscript *Measure theory for probabilists*. In particular, references to Chapters 1–5 are references to this manuscript.

The following books have guided me as references for the purpose of this manuscript.

- Durrett xxx
- Kallenberg, Olaf. Foundations of Modern Probability Theory. Springer, third edition, 2021
- Klenke, Achim. Probability theory. A comprehensive course. Springer, 2014

Throughout the manuscript, we will use a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ (recall from Definition 2.1). The integral with respect to \mathbf{P} is denoted by $\mathbf{E}[\cdot] := \mathbf{P}[\cdot]$ (recall from Chapter 3). Further, we abbreviate $\mathcal{L}^p := \mathcal{L}^p(\mathbf{P})$ if this does not lead to confusion (recall these spaces from Chapter 4).

Our aim in the present course is to provide the most important probabilistic statements available. Fundamental to this is the concept of the random variable, which we will examine in Chapter 6 (see also Definition 3.3). We will often consider the case of E -valued random variables, where E is a Polish space (see Appendix A in the lecture notes on measure theory). The most important theorems in stochastics are the strong law of large numbers (LLN, Theorem 8.21) and the central limit theorem (CLT, Theorem 10.8). These two theorems are limit statements for random variables, whereby it is important to note that the type of convergence in both theorems is fundamentally different. While the strong law of large numbers describes an almost sure convergence (refer to Remark 2.14), the central limit theorem is a statement about convergence in distribution (i.e. about the weak convergence of the distributions of the random variables; see Chapter 9). Consequently, one of the tasks will be to see the relationships between different types of convergence (see Chapter 7 and 9).

The present english version of this manuscript was written based on the German version with the help of DeepL.

Contents

6	Random variables	4
6.1	Repetition	4
6.2	Moments	8
6.3	Characteristic functions	9
7	Almost sure, stochastic and \mathcal{L}^p-convergence	12
7.1	Definition and examples	12
7.2	Almost sure convergence and convergence in probability	14
7.3	Convergence in probability and \mathcal{L}^p -convergence	15
8	Independence and the strong law	18
8.1	Definition and simple properties	18
8.2	Kolmogorov's 0-1 law	22
8.3	Sums of independent random variables	24
8.4	The Strong Law of Large Numbers	25
9	Weak convergence	30
9.1	Definition and simple properties	30
9.2	Prohorov' Theorem	36
9.3	Separating classes of functions	41
9.4	The theorem of Lévy	43
10	limit sets in distribution	47
10.1	Poisson convergence	47
10.2	The central limit theorem	50
10.3	Multidimensional limit sets	54
11	The conditional expectation	56
11.1	motivation	56
11.2	definition and properties	57
11.3	The case $\mathcal{G} = \sigma(X)$	61

6 Random variables

We usually use real-valued random variables $X : \Omega \rightarrow \mathbb{R}$ (i.e. Borel-measurable functions, i.e. random variables with values in \mathbb{R} , measurable with respect to the Borel σ -algebra in \mathbb{R} ; recall from Definition 1.7)). We will now recall several concepts from measure theory about random variables and will need directly in the following. We will mainly deal with connecting the lecture to measure theory on one side, and *Basic Probability* on the other side.

6.1 Repetition

Recall that we take a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ for granted. In Definition 3.5, we already defined several notions with a relationship to random variables, which we recall for completeness.

Remark 6.1 (Random variables and their distribution). *Let (Ω', \mathcal{F}') be a measurable space.*

1. *Every \mathcal{F}/\mathcal{F}' -measurable function X is called $(\Omega'$ -valued) random variable. If $(\Omega', \mathcal{F}') = (\mathbb{R}, \mathcal{B}(\mathbb{R}))$, it is called real-valued. The σ -algebra $\sigma(X) = \{X^{-1}(B) : B \in \mathcal{F}'\}$ is the X generated by σ -algebra (see Definition 3.3).*
2. *The probability measure $X_*\mathbf{P}$ on \mathcal{F}' (i.e. the image measure of \mathbf{P} under X ; see Section 2.5) is called distribution of X . Furthermore, if Y is a random variable and $X_*\mathbf{P} = Y_*\mathbf{P}$ (i.e. $\mathbf{P}(X \in A') = \mathbf{P}(Y \in A')$ for all $A' \in \mathcal{F}'$), then X and Y are identically distributed and we write $X \stackrel{d}{=} Y$. However, this notation should be used with caution, as the equality $X \stackrel{d}{=} Y$ cannot be achieved by equivalence transformations to other statements. (For example $X - Y \stackrel{d}{=} 0$ generally does not apply if X and Y are identically distributed).*
3. *For a family $(X_i)_{i \in I}$ of random variables, their joint distribution is given by $((X_i)_{i \in I})_*\mathbf{P}$. (This is the image measure under the mapping $(X_i)_{i \in I} : \omega \mapsto (X_i(\omega))_{i \in I}$).*
4. *We will use the following notation: If X is a random variable with distribution $N(\mu, \sigma^2)$. (This means means that $X : \Omega \mapsto \mathbb{R}$ is a measurable mapping and $X_*\mathbf{P} = \mu_{N(\mu, \sigma^2)}$; see example 2.22.) Then, we write $X \sim N(\mu, \sigma^2)$. Here, read ' \sim ' as has distribution.*
5. *Let λ be another measure on \mathcal{F} and $f : \Omega \rightarrow \mathbb{R}$ with $f \geq 0$ almost everywhere and $\lambda[f] = 1$. Then, X has the density f with respect to μ if and only if $X_*\mathbf{P} = f \cdot \lambda$ (see definition ??). Then, for $A \in \mathcal{F}$,*

$$\mathbf{P}(X \in A) = \mu[f, A].$$

In this case, the following applies for $g : \overline{\mathbb{R}} \rightarrow \overline{\mathbb{R}}$ that (see lemma 4.14)

$$\mathbf{E}[g(X)] = (X_*\mathbf{P})[g] = (f \cdot \mu)[g] = \mu[fg],$$

if the right-hand side exists.

6. *The monotonicity and linearity of the integral now means for random variables $X, Y \in \mathcal{L}^1$ and $a, b \in \mathbb{R}$:*

$$\begin{aligned} X \leq Y \text{ almost certainly} &\implies \mathbf{E}[X] \leq \mathbf{E}[Y], \\ \mathbf{E}[aX + bY] &= a\mathbf{E}[X] + b\mathbf{E}[Y]. \end{aligned}$$

Furthermore, according to Proposition 3.21,

$$\mathbf{E}[X] < \infty \implies \mathbf{P}(X < \infty) = 1.$$

Although the σ -algebra \mathcal{F} is given, in further sections, especially in the introduction of the conditional expectation in Chapter 11, the σ -algebra generated by X will play a special role. Simply put, a real-valued random variable Y is $\sigma(X)$ -measurable if and only if $Y = \varphi(X)$ for a Borel-measurable mapping φ . In other words, this means that the value of $Y(\omega)$ is known if you know $X(\omega)$, although you do not know what value ω has assumed. See also Exercise 3.38.

Lemma 6.2 (Measurability with respect to $\sigma(\mathbf{X})$). *Let (Ω', \mathcal{F}') be a measurable space and X a random variable with values in Ω' , and $Z : \Omega \rightarrow \overline{\mathbb{R}}$. Then, Z is $\sigma(X)$ -measurable if and only if there is a $\mathcal{F}'/\mathcal{B}(\overline{\mathbb{R}})$ -measurable mapping $\varphi : \Omega' \rightarrow \overline{\mathbb{R}}$ with $\varphi \circ X = Z$.*

Proof. ' \Leftarrow ': clear

' \Rightarrow ': It suffices to consider the case $Z \geq 0$; otherwise, we write $Z = Z^+ - Z^-$. First, let $Z = 1_A$ for $A \in \sigma(X)$. Then there is an $A' \in \mathcal{F}'$ with $X^{-1}(A') = A$, i.e. $Z = 1_{X^{-1}(A')} = 1_{A'} \circ X$, i.e. $\varphi = 1_{A'}$ fulfills the statement. Due to linearity, the statement is also true for simple functions, i.e. finite linear combinations of indicator functions. In the general case, there are simple functions $Z_1, Z_2, \dots \geq 0$ with $Z_n \uparrow Z$. In addition, there are \mathcal{F}' -measurable functions φ_n with $Z_n = \varphi_n \circ X$. Then $\varphi = \sup_n \varphi_n$ is again \mathcal{F}' -measurable and, since $Z \geq 0$, and

$$\varphi \circ X = \left(\sup_n \varphi_n\right) \circ X = \sup_n (\varphi_n \circ X) = \sup_n Z_n = Z.$$

□

We now briefly repeat the convergence theorems for integrals in the context of random variables.

Proposition 6.3 (Integral convergence theorems). *Let X, X_1, X_2, \dots be real-valued random variables.*

1. Lemma of Fatou, Theorem 3.27: *If $X_1, X_2, \dots \geq 0$, then*

$$\liminf_{n \rightarrow \infty} \mathbf{E}[X_n] \geq \mathbf{E}[\liminf_{n \rightarrow \infty} X_n].$$

2. Monotone convergence, Theorem 3.26: *If $X_1, X_2, \dots \in \mathcal{L}^1$ and $X_n \uparrow X$ is almost certain, then*

$$\lim_{n \rightarrow \infty} \mathbf{E}[X_n] = \mathbf{E}[X],$$

where both sides can take the value ∞ .

3. Dominated convergence, Theorem 3.28: *Let $X_n \xrightarrow{n \rightarrow \infty} X$ be almost certain and Y another real-valued random variable with $|X_1|, |X_2|, \dots \leq Y$ almost certain and $\mathbf{E}[Y] < \infty$. Then,*

$$\mathbf{E}[X_n] \xrightarrow{n \rightarrow \infty} \mathbf{E}[X].$$

We now collect (and re-prove) already known inequalities. They often help to estimate probabilities or expected values. Most of the inequalities are already known from the lecture on *Basic Probability*.

Proposition 6.4 (Markov and Chebyshev inequality). *1. Let X be a random variable with values in $\overline{\mathbb{R}}_+$ and $x \in \mathbb{R}_+$. Then the Markov inequality holds, i.e.,*

$$\mathbf{P}(X \geq x) \leq \frac{\mathbf{E}[X]}{x}.$$

2. *If X is a real-valued random variable and $p, x \in \mathbb{R}_+$, then the Chebyshev inequality holds, i.e.*

$$\mathbf{P}(|X| \geq x) \leq \frac{\mathbf{E}[|X|^p]}{x^p}.$$

Proof. 1. Since X is non-negative, we find $x \cdot 1_{X \geq x} \leq X$. Thus,

$$x \cdot \mathbf{P}(X \geq x) = \mathbf{E}[x \cdot 1_{X \geq x}] \leq \mathbf{E}[X],$$

and the inequality follows. The inequality in 2. follows from 1. by

$$\mathbf{P}(|X| \geq x) = \mathbf{P}(|X|^p \geq x^p) \leq \frac{\mathbf{E}[|X|^p]}{x^p}.$$

□

The next is Proposition 4.2.

Proposition 6.5 (Minkowski and Hölder inequalities). *Let X, Y be real-valued random variables.*

1. If $0 < p, q, r \leq \infty$ such that $\frac{1}{p} + \frac{1}{q} = \frac{1}{r}$. Then,

$$\mathbf{E}[|XY|^r]^{1/r} \leq \mathbf{E}[|X|^p]^{1/p} \cdot \mathbf{E}[|Y|^q]^{1/q} \quad (\text{H\"older inequality}) \quad (6.1)$$

Specifically, if $p = q = 2$

$$\mathbf{E}[|XY|] \leq \mathbf{E}[|X|^2]^{1/2} \cdot \mathbf{E}[|Y|^2]^{1/2}. \quad (\text{Cauchy-Schwartz inequality}) \quad (6.2)$$

2. For $1 \leq p \leq \infty$,

$$\mathbf{E}[|X + Y|^p]^{1/p} \leq \mathbf{E}[|X|^p]^{1/p} + \mathbf{E}[|Y|^p]^{1/p}, \quad 1 \leq p \leq \infty \quad (\text{Minkowski inequality}) \quad (6.3)$$

Proposition 6.6 (Jensen's inequality). Let $I \subseteq \mathbb{R}$ be an open interval and $X \in \mathcal{L}^1$ with values in I and $\varphi : I \rightarrow \mathbb{R}$ convex.¹ Then,

$$\mathbf{E}[\varphi(X)] \geq \varphi(\mathbf{E}[X]).$$

Proof. Since φ is convex, φ is continuous and

$$t \mapsto \frac{\varphi(tx + (1-t)y) - \varphi(y)}{t(x-y)}$$

is monotonically decreasing for $y \leq x$. In particular, for $y \in I$ there exists

$$\lambda(y) := \lim_{x \downarrow y} \frac{\varphi(x) - \varphi(y)}{x - y} = \lim_{t \downarrow 0} \frac{\varphi(tx + (1-t)y) - \varphi(y)}{t(x-y)} \quad (6.4)$$

and

$$\frac{\varphi(x) - \varphi(y)}{x - y} \geq \lambda(y) \implies \varphi(y) + \lambda(y)(x - y) \leq \varphi(x) \quad (6.5)$$

for all $x \in I$. (For $y > x$ one argues analogously as above).

Note, since I is an interval, we have $\mathbf{E}[X] \in I$. According to (6.5), for $x \in I$ with $y = \mathbf{E}[X]$

$$\varphi(x) \geq \varphi(\mathbf{E}[X]) + \lambda(\mathbf{E}[X])(x - \mathbf{E}[X])$$

and thus

$$\mathbf{E}[\varphi(X)] \geq \varphi(\mathbf{E}[X]) + \lambda(\mathbf{E}[X])\mathbf{E}[X - \mathbf{E}[X]] = \varphi(\mathbf{E}[X]). \quad \square$$

Jensen's inequality can be used to show, for example, that $\mathcal{L}^q \subseteq \mathcal{L}^p$ for $p \leq q$. Alternatively, you can read this property from Proposition 4.3.

Lemma 6.7 (p -fold and q -fold integrable random variable). Let $q > 0$ and $X \in \mathcal{L}^q$ be a real-valued random variable. Then for $p \leq q$

$$\mathbf{E}[|X|^p] \leq \mathbf{E}[|X|^q]^{p/q}.$$

In particular, $\mathcal{L}^q \subseteq \mathcal{L}^p$.

Proof. The mapping $y \mapsto y^{p/q}$ is concave on \mathbb{R}_+ , so with Jensen's inequality,

$$\mathbf{E}[|X|^p] = \mathbf{E}[(|X|^q)^{p/q}] \leq \mathbf{E}[|X|^q]^{p/q}. \quad \square$$

¹A mapping $\varphi : I \rightarrow \mathbb{R}$ is convex if $\varphi(tx + (1-t)y) \leq t\varphi(x) + (1-t)\varphi(y)$ for all $0 \leq t \leq 1$ and $x, y \in I$.

6.2 Moments

From the lecture *Basic Probability*, we know terms such as expected value, variance and covariance. When we repeat them now, you will see that all calculation rules that you learned also apply in the measure theoretic sense. The only difference in the formulation is that $\mathbf{E}[\cdot]$ is the integral with respect to a probability measure.

Definition 6.8 (Moments). *Let X, Y be real-valued random variables. Then, if it exists, $\mathbf{E}[X]$ is the expected value of X . Furthermore, if it exists, the variance of X is given by*

$$\mathbf{V}[X] := \mathbf{E}[(X - \mathbf{E}[X])^2].$$

If it exists, the covariance of X and Y is given by

$$\mathbf{COV}[X, Y] := \mathbf{E}[(X - \mathbf{E}[X])(Y - \mathbf{E}[Y])].$$

If $\mathbf{COV}[X, Y] = 0$, we say that X and Y are uncorrelated. Furthermore, $\mathbf{E}[X^p]$ for $p > 0$ is the p -th moment of X and $\mathbf{E}[(X - \mathbf{E}[X])^p]$ is the centered p -th moment of X .

We recall a few properties here.

Proposition 6.9 (Properties of the second moments). *Let $X, Y \in \mathcal{L}^2$ be real-valued random variables. Then, $\mathbf{V}[X], \mathbf{V}[Y], |\mathbf{COV}[X, Y]| < \infty$ and*

$$\begin{aligned} \mathbf{V}[X] &= \mathbf{E}[X^2] - (\mathbf{E}[X])^2, \\ \mathbf{COV}[X, Y] &= \mathbf{E}[XY] - \mathbf{E}[X] \cdot \mathbf{E}[Y]. \end{aligned}$$

The Cauchy-Schwartz inequality holds, i.e.

$$\mathbf{COV}[X, Y]^2 \leq \mathbf{V}[X] \cdot \mathbf{V}[Y].$$

If $X_1, \dots, X_n \in \mathcal{L}^2$, the identity of Bienamyé holds, i.e.

$$\mathbf{V}\left[\sum_{k=1}^n X_k\right] = \sum_{k=1}^n \mathbf{V}[X_k] + 2 \sum_{1 \leq k < l \leq n} \mathbf{COV}[X_k, X_l].$$

Proof. For the first statement, since $\mathbf{V}[X] = \mathbf{COV}[X, X]$ by definition, it is sufficient to show the second equation. This equation follows from the linearity of the expected value by means of

$$\begin{aligned} \mathbf{COV}[X, Y] &= \mathbf{E}[(X - \mathbf{E}[X])(Y - \mathbf{E}[Y])] \\ &= \mathbf{E}[XY] - \mathbf{E}[\mathbf{E}[X]Y] - \mathbf{E}[X\mathbf{E}[Y]] + \mathbf{E}[X]\mathbf{E}[Y] \\ &= \mathbf{E}[XY] - \mathbf{E}[X]\mathbf{E}[Y]. \end{aligned}$$

The Cauchy-Schwartz inequality follows by applying Proposition 6.5 using the random variables $X - \mathbf{E}[X]$ and $Y - \mathbf{E}[Y]$. In particular, $|\mathbf{COV}[X, Y]| < \infty$. For the equation of Bienamyé let wlog $\mathbf{E}[X_k] = 0, k = 1, \dots, n$ (otherwise we use the random variables $X_k - \mathbf{E}[X_k]$). Then,

$$\begin{aligned} \mathbf{V}\left[\sum_{k=1}^n X_k\right] &= \mathbf{E}\left[\left(\sum_{k=1}^n X_k\right)^2\right] = \sum_{k=1}^n \sum_{l=1}^n \mathbf{E}[X_k X_l] = \sum_{k=1}^n \mathbf{E}[X_k^2] + 2 \sum_{1 \leq k < l \leq n} \mathbf{E}[X_k X_l] \\ &= \sum_{k=1}^n \mathbf{V}[X_k] + 2 \sum_{1 \leq k < l \leq n} \mathbf{COV}[X_k, X_l]. \end{aligned} \quad \square$$

Proposition 6.10 (Moments of non-negative random variables). *Let X be a random variable with values in \mathbb{R}_+ . Then,*

$$\mathbf{E}[X^p] = p \int_0^\infty \mathbf{P}(X > t)t^{p-1}dt = p \int_0^\infty \mathbf{P}(X \geq t)t^{p-1}dt.$$

Proof. Note that both, $\mathbf{E}[\cdot]$ and $\int \cdot dt$ are integrals. We use Fubini's theorem in order to be able to change the order of integration,

$$\mathbf{E}[X^p] = p\mathbf{E}\left[\int_0^X t^{p-1}dt\right] = p \int_0^\infty \mathbf{E}\left[1_{X>t}t^{p-1}\right]dt = p \int_0^\infty \mathbf{P}(X > t)t^{p-1}dt.$$

The proof of the second equation is analogous. □

6.3 Characteristic functions and Laplace transforms

We now introduce expected values of certain functions of random variables. This results in the characteristic function (of the distribution of real-valued random variables) and the Laplace transform (of the distribution of non-negative random variables). (Both are covered in some courses on Basic Probability Theory, but not in all.) These functions are useful since they allow the calculation of moments (see Proposition 6.14). In addition, later in Proposition 9.25, we will show that these functions uniquely determine the underlying measure.

Definition 6.11 (Characteristic function and Laplace transform).

1. The characteristic function of an \mathbb{R}^d -valued random variable X is given by

$$\psi_X := \psi_{X_*\mathbf{P}} := \begin{cases} \mathbb{R}^d & \rightarrow \mathbb{C}, \\ t & \mapsto \mathbf{E}[e^{itX}] := \mathbf{E}[\cos(tX)] + i\mathbf{E}[\sin(tX)], \end{cases}$$

where $tx := \langle t, x \rangle$ is the scalar product in \mathbb{R}^d .

2. The Laplace transform of X is given by

$$\mathcal{L}_X := \mathcal{L}_{X_*\mathbf{P}} := \begin{cases} \mathbb{R}^d & \rightarrow \mathbb{R}, \\ t & \mapsto \mathbf{E}[e^{-tX}], \end{cases}$$

provided the integral on the right-hand side exists. (The Laplace transform is used frequently for probability measures on \mathbb{R}_+^d .)

Proposition 6.12 (Properties of characteristic functions). *Let X, Y be random variables with values in \mathbb{R}^d and characteristic functions ψ_X, ψ_Y . Then,*

1. $|\psi_X(t)| \leq 1$ for each $t \in \mathbb{R}^d$ and $\psi_X(0) = 1$.
2. ψ_X is uniformly continuous.
3. $\psi_{aX+b}(t) = \psi_X(at)e^{ibt}$ for all $a \in \mathbb{R}, b \in \mathbb{R}^d$.

Proof. 1. is clear. For uniform continuity, we have the bound

$$\begin{aligned} |e^{ihx} - 1| &= \sqrt{|\cos(hx) + i\sin(hx) - 1|^2} = \sqrt{(\cos(hx) - 1)^2 + \sin^2(hx)} \\ &= \sqrt{2(1 - \cos(hx))} = 2|\sin(hx/2)| \leq |hx| \wedge 2. \end{aligned}$$

From this, 2. follows because of

$$\begin{aligned} \sup_{t \in \mathbb{R}^d} |\psi_X(t+h) - \psi_X(t)| &= \sup_{t \in \mathbb{R}^d} |\mathbf{E}[e^{i(t+h)X} - e^{itX}]| = \sup_{t \in \mathbb{R}^d} |\mathbf{E}[e^{itX}(e^{ihX} - 1)]| \\ &\leq \mathbf{E}[|e^{ihX} - 1|] \leq \mathbf{E}[|hX| \wedge 2] \xrightarrow{h \rightarrow 0} 0. \end{aligned}$$

For 3. we calculate using linearity

$$\mathbf{E}[e^{it(aX+b)}] = e^{itb} \mathbf{E}[e^{i(at)X}] = e^{itb} \psi_X(at). \quad \square$$

Example 6.13 (Examples of characteristic functions functions). 1. The characteristic function of $X \sim B(n, p)$ is given by

$$\psi_{B(n,p)}(t) = (1 - p + pe^{it})^n.$$

Indeed: By definition,

$$\mathbf{E}[e^{itX}] = \sum_{k=0}^n \binom{n}{k} p^k (1-p)^{n-k} e^{itk} = (1 - p + pe^{it})^n.$$

2. The characteristic function of $X \sim Poi(\gamma)$ is given by

$$\psi_{Poi(\gamma)}(t) = e^{\gamma(e^{it}-1)}.$$

Indeed: We calculate

$$\psi_{Poi(\gamma)}(t) = e^{-\gamma} \sum_{n=0}^{\infty} \frac{\gamma^n e^{itn}}{n!} = e^{\gamma(e^{it}-1)}.$$

3. The characteristic function of $X \sim N(\mu, \sigma^2)$ is given by

$$\psi_{N(\mu, \sigma^2)}(t) = e^{it\mu} e^{-\sigma^2 t^2/2}.$$

Indeed: We use that $X \sim \sigma Z + \mu$ for $Z \sim N(0, 1)$. According to Proposition 6.12.2, it is sufficient to compute this assertion for $\mu = 0, \sigma^2 = 1$. For this case, using partial integration,

$$\frac{d}{dt} \psi_{N(0,1)}(t) = \frac{i}{\sqrt{2\pi}} \int x e^{-x^2/2} e^{itx} dx = -\frac{i}{\sqrt{2\pi}} \int e^{-x^2/2} i t e^{itx} dx = -t \psi_{N(0,1)}(t).$$

This differential equation with $\psi_{N(0,1)}(0) = 1$ has the unique solution $\psi_{N(0,1)}(t) = e^{-t^2/2}$.

4. The Laplace transform of $X \sim \exp(\gamma)$ is given by

$$\mathcal{L}_{\exp(\gamma)}(t) = \frac{\gamma}{\gamma + t}.$$

Indeed: A straight-forward calculation reveals

$$\mathbf{E}[e^{-tX}] = \int_0^{\infty} \gamma e^{-\gamma x} e^{-tx} dx = \frac{\gamma}{\gamma + t}.$$

Characteristic functions and Laplace transforms are a simple tool to calculate the moments of random variables.

Proposition 6.14 (Characteristic function and moments). *Let X be a real-valued random variable.*

1. If $X \in \mathcal{L}^p$, then ψ_X is p -times continuously differentiable and for $k = 0, \dots, p$,

$$\psi_X^{(k)}(t) = \mathbf{E}[(iX)^k e^{itX}].$$

In particular, $\psi_X^{(k)}(0) = i^k \mathbf{E}[X^k]$.

2. In particular, if $X \in \mathcal{L}^2$, then

$$\psi_X(t) = 1 + it\mathbf{E}[X] - \frac{t^2}{2}\mathbf{E}[X^2] + \varepsilon(t)t^2$$

as $\varepsilon(t) \xrightarrow{t \rightarrow 0} 0$.

Proof. 1. With $|X|^p$ also $|X|^p \vee 1$ is integrable. Thus, since $|X|^k \leq |X|^p \vee 1$, all $|X|^k$ can be dominated by an integrable random variable and the right-hand side exists. Since the statement is obviously true for $k = 0$, we assume that it is valid for $k < n$. Then

$$\left| \frac{d^{k+1}}{dt^{k+1}} e^{itX} \right| = \lim_{h \rightarrow 0} \left| \frac{(iX)^k e^{i(t+h)X} - (iX)^k e^{itX}}{h} \right| \leq \left| X^k \frac{e^{ihX} - 1}{h} \right| \leq |X^{k+1}|.$$

Due to dominated convergence, the derivative and integral can be interchanged and it follows

$$\psi_X^{(k+1)}(t) = \mathbf{E} \left[\frac{d}{dt} (iX)^k e^{itX} \right] = \mathbf{E}[(iX)^{k+1} e^{itX}].$$

The continuity of the derivative also follows with dominated convergence.

2. For the estimation, we need the Taylor expansion of ψ_X with remainder term. We have

$$e^{itX} = 1 + itX - \frac{t^2 X^2}{2} (\cos(\theta_1 tX) + i \sin(\theta_2 tX))$$

with random numbers θ_1, θ_2 , so that $|\theta_1|, |\theta_2| \leq 1$. Therefore we get

$$\psi_X(t) = 1 + it\mathbf{E}[X] - \frac{t^2}{2}\mathbf{E}[X^2] + \varepsilon(t)t^2$$

with $2\varepsilon(t) = \mathbf{E}[X^2(1 - \cos(\theta_1 tX) + i \sin(\theta_2 tX))] \xrightarrow{t \rightarrow 0} 0$ from dominated convergence. \square

Example 6.15 (Moments of the exponential and normal distribution). 1. Let $X \sim \exp(\gamma)$.

We know that $\mathcal{L}_{\exp(\gamma)}(t) = \gamma/(\gamma+t)$ from Example 6.13.4. From this and the last Proposition,

$$\mathbf{E}[X^n] = (-1)^n \frac{d^n}{dt^n} \mathbf{E}[e^{-tX}] \Big|_{t=0} = (-1)^n \frac{d^n}{dt^n} \frac{\gamma}{\gamma+t} \Big|_{t=0} = \frac{n! \gamma}{(\gamma+t)^{n+1}} \Big|_{t=0} = \frac{n!}{\gamma^n}.$$

2. For $X \sim N(\mu, \sigma^2)$, we already know $\psi_{N(\mu, \sigma^2)}(t) = e^{it\mu - \sigma^2 t^2 / 2}$. For small t we develop this with

$$\psi_{N(\mu, \sigma^2)}(t) = 1 + it\mu - \sigma^2 t^2 / 2 - \mu^2 t^2 / 2 + \varepsilon(t)t^2$$


with $\varepsilon(t) \xrightarrow{t \rightarrow 0} 0$. From this one reads by means of Proposition 6.14.2 that

$$\mathbf{E}[X] = \mu, \quad \mathbf{V}[X] = \mathbf{E}[X^2] - \mu^2 = \sigma^2.$$

7 Almost sure, stochastic and \mathcal{L}^p -convergence

It is already known from lectures on *Analysis* that there are different types of convergence, such as uniform and pointwise convergence. We will now discuss the most important types of convergence with respect to random variables. (For definitions, see below.)

In our course on measure theory, we have already seen almost sure convergence. In addition, we will discuss convergence in probability and the \mathcal{L}^p -convergence (see also Section 4). In Section 9, we will learn about convergence in distribution (which is the same as the weak convergence of the distributions of random variables). The following diagram summarizes all types of convergence:



7.1 Definition and examples

Let's start with some definitions.

Definition 7.1 (Almost sure convergence and convergence in probability). *Let X, X_1, X_2, \dots be random variables with values in a metric space (E, r) .*

1. If

$$\mathbf{P}\left(\lim_{n \rightarrow \infty} r(X_n, X) = 0\right) = 1,$$

we say that the sequence X_1, X_2, \dots converges almost surely to X and write $X_n \xrightarrow{n \rightarrow \infty}_{as} X$.

2. If, for all $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \mathbf{P}(r(X_n, X) > \varepsilon) = 0,$$

we say that the sequence X_1, X_2, \dots converges to X in probability (or stochastically) and write $X_n \xrightarrow{n \rightarrow \infty}_p X$.

3. If the random variables are real-valued and for some $p > 0$

$$\lim_{n \rightarrow \infty} \mathbf{E}[|X_n - X|^p] = 0,$$

we say that the sequence X_1, X_2, \dots converges in \mathcal{L}^p (or in the p -th mean) to X and also write $X_n \xrightarrow{n \rightarrow \infty}_{\mathcal{L}^p} X$.

Remark 7.2 (Properties of \mathcal{L}^p convergence). *From section 4 we already know a lot about the \mathcal{L}^p -convergence. For example, if X, X_1, X_2, \dots is such that $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^q X$ and $p < q$, then $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$ according to Proposition 4.7. In addition, the spaces \mathcal{L}^p are complete according to Proposition 4.8. So, if for all $\varepsilon > 0$, there is an $N \in \mathbb{N}$ such that for all $m, n \geq n$*

$$\mathbf{E}[|X_n - X_m|^p] < \varepsilon,$$

then there is a random variable $X \in \mathcal{L}^p$ with $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$.

Example 7.3 (Counterexamples). *If we look at the diagram at the beginning of the chapter, we can see that convergence in probability follows from almost sure convergence, but not vice versa. Furthermore, convergence in probability follows from \mathcal{L}^1 convergence, but not even almost sure convergence implies \mathcal{L}^1 convergence. We first give two examples for these two cases.*

1. *Convergence in probability does not imply almost sure convergence: Let U be a $[0, 1]$ uniformly distributed random variable. Further we set*

$$\begin{aligned} A_1 &= [0, \frac{1}{2}], & A_2 &= [\frac{1}{2}, 1], \\ A_3 &= [0, \frac{1}{4}], & A_4 &= [\frac{1}{4}, \frac{2}{4}], & A_5 &= [\frac{2}{4}, \frac{3}{4}], & A_6 &= [\frac{3}{4}, 1], \\ &\dots & & & & & & \end{aligned}$$

and $X_n := 1_{U \in A_n}$. Then it is clear for $0 < \varepsilon < 1$

$$\lim_{n \rightarrow \infty} \mathbf{P}(|X_n| > \varepsilon) = \lim_{n \rightarrow \infty} \mathbf{P}(U \in A_n) = 0,$$

i.e. $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p 0$, but for each $n \in \mathbb{N}$ there is an $m > n$ with $X_m = 1$. Therefore, X_1, X_2, \dots does not converge almost surely to 0.

2. *Almost sure convergence does not imply \mathcal{L}^1 convergence: Let U again be a uniformly distributed random variable on $[0, 1]$ random variable. Further, $B_n = [0, \frac{1}{n}]$ and $Y_n = n \cdot 1_{U \in B_n}$. Then $Y_n \xrightarrow{n \rightarrow \infty} \text{fs } Y = \infty \cdot 1_{U=0}$, so $Y = 0$ is almost certain. On the other hand*

$$\mathbf{E}[Y_n] = n \cdot \mathbf{P}(U \in A_n) = 1,$$

so Y_1, Y_2, \dots does not converge to 0 in \mathcal{L}^1 .

Lemma 7.4 (Limit in probability is (almost surely) unique). *Let X, Y, X_1, X_2, \dots be random variables with values in a metric space space (E, r) and $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$, as well as $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p Y$. Then $X = Y$ almost surely.*

Proof. For all $\varepsilon > 0$,

$$\begin{aligned} \mathbf{P}(r(X, Y) > 2\varepsilon) &\leq \mathbf{P}(r(X_n, X) > \varepsilon \text{ or } r(X_n, Y) > \varepsilon) \\ &\leq \mathbf{P}(r(X_n, X) > \varepsilon) + \mathbf{P}(r(X_n, Y) > \varepsilon) \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Therefore,

$$\mathbf{P}(X \neq Y) = \mathbf{P}\left(\bigcup_{k=1}^{\infty} \{r(X, Y) > 1/k\}\right) \leq \sum_{k=1}^{\infty} \mathbf{P}(r(X, Y) > 1/k) = 0,$$

and the statement follows. □

7.2 Almost sure convergence and convergence in probability

We now show a result that relates almost sure convergence and convergence in probability.

Lemma 7.5 (Characterization of convergence in probability). *Let X, X_1, X_2, \dots be random variables with values in a metric space (E, r) . Then,*

$$X_n \xrightarrow[n \rightarrow \infty]{p} X \iff \mathbf{E}[r(X_n, X) \wedge 1] \xrightarrow[n \rightarrow \infty]{} 0. \quad (7.1)$$

Proof. If $X_n \xrightarrow[n \rightarrow \infty]{p} X$, then for all $\varepsilon > 0$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbf{E}[r(X_n, X) \wedge 1] &= \lim_{n \rightarrow \infty} \mathbf{E}[r(X_n, X) \wedge 1, r(X_n, X) \leq \varepsilon] + \mathbf{E}[r(X_n, X) \wedge 1, r(X_n, X) > \varepsilon] \\ &\leq \lim_{n \rightarrow \infty} (\varepsilon + \mathbf{P}(r(X_n, X) > \varepsilon)) = \varepsilon, \end{aligned}$$

which shows the right-hand side since $\varepsilon > 0$ was arbitrary. However, if the right-hand side applies, the Chebyshev inequality for $0 < \varepsilon \leq 1$ implies that

$$\mathbf{P}(r(X_n, X) > \varepsilon) \leq \frac{\mathbf{E}[r(X_n, X) \wedge 1]}{\varepsilon} \xrightarrow[n \rightarrow \infty]{} 0. \quad \square$$

Proposition 7.6 (Convergence in probability and almost sure convergence). *Let X, X_1, X_2, \dots be random variables with values in a metric space (E, r) . Then, the following are equivalent:*

1. $X_n \xrightarrow[n \rightarrow \infty]{p} X$
2. For each sequence $(n_k)_{k=1,2,\dots}$ there is a subsequence $(n_{k_\ell})_{\ell=1,2,\dots}$ with $X_{n_{k_\ell}} \xrightarrow[\ell \rightarrow \infty]{as} X$.

In particular,

$$X_n \xrightarrow[n \rightarrow \infty]{fs} X \implies X_n \xrightarrow[n \rightarrow \infty]{p} X.$$

Proof. 1. \rightarrow 2.: Because of (7.1), for each sequence $(n_k)_{k=1,2,\dots}$, there is a subsequence $(n_{k_\ell})_{\ell=1,2,\dots}$ such that (using monotone convergence in the first equality)

$$\mathbf{E} \left[\sum_{\ell=1}^{\infty} (r(X_{n_{k_\ell}}, X) \wedge 1) \right] = \sum_{\ell=1}^{\infty} \mathbf{E}[r(X_{n_{k_\ell}}, X) \wedge 1] < \infty.$$

This means that

$$1 = \mathbf{P} \left(\sum_{\ell=1}^{\infty} (r(X_{n_{k_\ell}}, X) \wedge 1) < \infty \right) \leq \mathbf{P} \left(\limsup_{\ell \rightarrow \infty} r(X_{n_{k_\ell}}, X) = 0 \right) \leq 1,$$

i.e. $X_{n_{k_\ell}} \xrightarrow[\ell \rightarrow \infty]{fs} X$.

2. \rightarrow 1.: Let's assume that 1. is not valid. (We must show that 2. cannot hold.) Because of (7.1), there is $\varepsilon > 0$ and a subsequence $(n_k)_{k=1,2,\dots}$, so that $\lim_{k \rightarrow \infty} \mathbf{E}[r(X_{n_k}, X) \wedge 1] > \varepsilon$. Assuming that there is a subsequence $(n_{k_\ell})_{\ell=1,2,\dots}$ such that $X_{n_{k_\ell}} \xrightarrow[\ell \rightarrow \infty]{as} X$. Then also

$$\lim_{\ell \rightarrow \infty} \mathbf{E}[r(X_{n_{k_\ell}}, X) \wedge 1] = \mathbf{E} \left[\lim_{\ell \rightarrow \infty} r(X_{n_{k_\ell}}, X) \wedge 1 \right] = 0$$

due to dominated convergence, i.e. a contradiction. So we have found a sequence $(n_k)_{k=1,2,\dots}$ for which there is no further subsequence $(n_{k_\ell})_{\ell=1,2,\dots}$ with $X_{n_{k_\ell}} \xrightarrow[\ell \rightarrow \infty]{as} X$, and we have shown that 2. does not hold. \square

7.3 Convergence in probability and \mathcal{L}^p -convergence

In Example 7.3 we had already seen that almost sure convergence (as well as convergence in probability) does not imply \mathcal{L}^1 -convergence. This is not surprising, since the theorem of dominated convergence states that a sequence X_1, X_2, \dots , which converges almost surely to X and has an integrable dominating random variable converges in \mathcal{L}^1 to X . If the almost sure convergence implies the \mathcal{L}^1 convergence, one would not need to make the requirement of an integrable dominating random variable. In the following, we want to find conditions of the integrable dominating random variable in order to suffice for \mathcal{L}^1 convergence. See theorem 7.11 and Corollary 7.12. The concept of uniform integrability is central to this, see Definition 7.7.

Definition 7.7 (Uniform integrability). *A family $(X_i)_{i \in I}$ is called uniformly integrable if*

$$\inf_K \sup_{i \in I} \mathbf{E}[|X_i|; |X_i| > K] = 0.$$

Example 7.8 (Uniform integrability). *1. Let $Y \in \mathcal{L}^1$ and $(X_i)_{i \in I}$ with $\sup_i |X_i| \leq |Y|$. Then $(X_i)_{i \in I}$ is uniformly integrable since*

$$\sup_{i \in I} \mathbf{E}[|X_i|; |X_i| > K] \leq \mathbf{E}[|Y|; |Y| > K] \xrightarrow{K \rightarrow \infty} 0$$

by dominated convergence. In particular, every $Y \in \mathcal{L}^1$ is uniformly integrable.

- 2. Every finite family $(X_i)_{i=1, \dots, n}$ with $X_1, \dots, X_n \in \mathcal{L}^1$ is uniformly integrable, because $\sup_{1 \leq i \leq n} |X_i| \in \mathcal{L}^1$ and therefore, 1. applies with $Y = \sup_{1 \leq i \leq n} |X_i|$.*
- 3. Let us consider the example 7.3.2 Here, for $n > K$*

$$\mathbf{E}[|Y_n|; |Y_n| > K] = \mathbf{E}[Y_n] = 1.$$

In particular, $(Y_n)_{n=1, 2, \dots}$ is not uniformly integrable.

- 4. Let $p > 1$. Then $(X_i)_{i \in I}$ with $X_i \in \mathcal{L}^p, i \in I$ is uniformly integrable if $\sup_{i \in I} \|X_i\|_p < \infty$. This is because $K^{p-1}|X_i|1_{|X_i| > K} \leq |X_i|^p$, therefore*

$$\sup_{i \in I} \mathbf{E}[|X_i|; |X_i| > K] \leq \sup_{i \in I} \frac{\mathbf{E}[|X_i|^p]}{K^{p-1}} \xrightarrow{K \rightarrow \infty} 0.$$

Lemma 7.9 (Characterization of uniform integrability). *Let $(X_i)_{i \in I}$ be a family of random variables. Then, the following are equivalent:*

- 1. $(X_i)_{i \in I}$ is uniformly integrable.*
- 2. It holds*

$$\sup_{i \in I} \mathbf{E}[|X_i|] < \infty \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \sup_{A: \mathbf{P}(A) < \varepsilon} \sup_{i \in I} \mathbf{E}[|X_i|; A] = 0.$$

- 3. It holds*

$$\lim_{K \rightarrow \infty} \sup_{i \in I} \mathbf{E}[(|X_i| - K)^+] = 0.$$

4. There exists a function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $\frac{f(x)}{x} \xrightarrow{x \rightarrow \infty} \infty$ and $\sup_{i \in I} \mathbf{E}[f(|X_i|)] < \infty$.

If one of the four statements is true, the function f in 4. can be chosen to be monotonically increasing and convex.

Proof. '1. \rightarrow 2.': Let $\delta > 0$ be given and $K = K_\delta$ such that $\sup_{i \in I} \mathbf{E}[|X_i|; |X_i| > K] \leq \delta$. Then, for $A \in \mathcal{F}$,

$$\mathbf{E}[|X_i|; A] = \mathbf{E}[|X_i|; A \cap \{|X_i| > K\}] + \mathbf{E}[|X_i|; A \cap \{|X_i| \leq K\}] \leq \delta + K \cdot \mathbf{P}(A).$$

In particular,

$$\sup_{i \in I} \mathbf{E}[|X_i|] = \sup_{i \in I} \mathbf{E}[|X_i|; \Omega] \leq \delta + K < \infty$$

and

$$\sup_{A: \mathbf{P}(A) < \varepsilon} \sup_{i \in I} \mathbf{E}[|X_i|; A] \leq \delta + K\varepsilon \xrightarrow{\varepsilon \rightarrow 0} \delta.$$

Since $\delta > 0$ was arbitrary,

$$\lim_{\varepsilon \rightarrow 0} \sup_{A: \mathbf{P}(A) < \varepsilon} \sup_{i \in I} \mathbf{E}[|X_i|; A] = 0.$$

'2. \Rightarrow 3.': First, we note that $(|X_i| - K)^+ \leq |X_i| 1_{|X_i| \geq K}$. Let $\varepsilon > 0$. Choose $K = K_\varepsilon$ large enough so that – using the Markov inequality –

$$\sup_{i \in I} \mathbf{P}(|X_i| > K) \leq \sup_{i \in I} \frac{\mathbf{E}[|X_i|]}{K} < \varepsilon.$$

This means that 3. follows from

$$\begin{aligned} \lim_{K \rightarrow \infty} \sup_{i \in I} \mathbf{E}[(|X_i| - K)^+] &= \lim_{\varepsilon \rightarrow 0} \sup_{i \in I} \mathbf{E}[(|X_i| - K_\varepsilon)^+] \leq \lim_{\varepsilon \rightarrow 0} \sup_{i \in I} \mathbf{E}[|X_i|; |X_i| > K_\varepsilon] \\ &\leq \lim_{\varepsilon \rightarrow 0} \sup_{A: \mathbf{P}(A) < \varepsilon} \sup_{i \in I} \mathbf{E}[|X_i|; A] = 0. \end{aligned}$$

'3. \Rightarrow 4.': There is a sequence K_1, K_2, \dots with $K_n \uparrow \infty$ and $\sup_{i \in I} \mathbf{E}[(|X_i| - K_n)^+] \leq 2^{-n}$. We set

$$f(x) := \sum_{n=1}^{\infty} (x - K_n)^+.$$

Then f is monotonically increasing and convex as a sum of convex functions. Furthermore, for $x \geq 2K_n$,

$$\frac{f(x)}{x} \geq \sum_{k=1}^n \left(1 - \frac{K_k}{x}\right) \geq \frac{n}{2},$$

thus $\frac{f(x)}{x} \xrightarrow{x \rightarrow \infty} \infty$. Because of monotone convergence,

$$\mathbf{E}[f(|X_i|)] = \sum_{n=1}^{\infty} \mathbf{E}[(|X_i| - K_n)^+] \leq \sum_{n=1}^{\infty} 2^{-n} = 1.$$

'4. \Rightarrow 1.': Set $a_K := \inf_{x \geq K} \frac{f(x)}{x}$, so that $a_K \xrightarrow{K \rightarrow \infty} \infty$. Thus,

$$\sup_{i \in I} \mathbf{E}[|X_i|; |X_i| \geq K] \leq \frac{1}{a_K} \sup_{i \in I} \mathbf{E}[f(|X_i|); |X_i| \geq K] \leq \frac{1}{a_K} \sup_{i \in I} \mathbf{E}[f(|X_i|)] \xrightarrow{K \rightarrow \infty} 0. \quad \square$$

Example 7.10 (Differences and uniform integrability). For $X \in \mathcal{L}^1$, $(X_i)_{i \in I}$ is uniformly integrable iff $(X_i - X)_{i \in I}$ is uniformly integrable.

To see this, let $(X_i)_{i \in I}$ be uniformly integrable. According to Example 7.8.2, X is uniformly integrable. Furthermore,

$$\sup_{i \in I} \mathbf{E}[|X_i - X|] \leq \mathbf{E}[|X|] + \sup_{i \in I} \mathbf{E}[|X_i|] < \infty$$

and

$$\sup_{A: \mathbf{P}(A) < \varepsilon} \sup_{i \in I} \mathbf{E}[|X_i - X|; A] \leq \sup_{A: \mathbf{P}(A) < \varepsilon} \sup_{i \in I} \mathbf{E}[|X_i|; A] + \sup_{A: \mathbf{P}(A) < \varepsilon} \mathbf{E}[|X|; A] \xrightarrow{\varepsilon \rightarrow 0} 0,$$

i.e. according to Lemma 7.9, $(X_i - X)_{i \in I}$ is uniformly integrable. The inverse follows analogously.

Theorem 7.11 (Convergence in probability and convergence in the p -th mean). Let X_1, X_2, \dots be a sequence in \mathcal{L}^p with $1 \leq p < \infty$. The following statements are equivalent:

1. There is a measurable function $X \in \mathcal{L}^p$ with $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$.
2. The family $(|X_i|^p)_{i=1,2,\dots}$ is uniformly integrable and there is a measurable function X with $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$.

If 1. or 2. applies, then the limits coincide almost surely.

Proof. 1. \rightarrow 2.: First, due to Chebyshev's inequality for every $\varepsilon > 0$

$$\mathbf{P}(|X_n - X| > \varepsilon) \leq \frac{\mathbf{E}[|X_n - X|^p]}{\varepsilon^p} = \frac{\|X_n - X\|_p^p}{\varepsilon^p} \xrightarrow{n \rightarrow \infty} 0,$$

i.e. convergence in probability applies. For the proof of uniform integrability, we use Lemma 7.9. Let $\varepsilon > 0$ and $N = N_\varepsilon$ such that $\|X_n - X\|_p < \varepsilon$ for $n \geq N$. Then for $A \in \mathcal{F}$, with Minkowski's inequality,

$$\begin{aligned} \sup_{n \in \mathbb{N}} (\mathbf{E}[|X_n|^p; A])^{1/p} &= \sup_{n \in \mathbb{N}} \|X_n 1_A\|_p \\ &\leq \sup_{n < N} \|X_n 1_A\|_p + \sup_{n \geq N} \|(X_n - X) 1_A\|_p + \|X 1_A\|_p \\ &\leq \sup_{n < N} (\mathbf{E}[|X_n|^p; A])^{1/p} + \varepsilon + (\mathbf{E}[|X|^p; A])^{1/p}. \end{aligned}$$

Using $A = \Omega$, we find $\sup_{n \in \mathbb{N}} (\mathbf{E}[|X_n|^p]) < \infty$. Moreover, since N is finite, we find

$$\lim_{\delta \rightarrow 0} \sup_{A: \mathbf{P}(A) < \delta} \sup_{n \in \mathbb{N}} \mathbf{E}[|X_n|^p; A] \leq \varepsilon^p.$$

Because $\varepsilon > 0$ was arbitrary, the assertion follows.

2. \rightarrow 1.: Since $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$, according to proposition 7.6 there is a subsequence n_1, n_2, \dots with $X_{n_k} \xrightarrow{k \rightarrow \infty} X$ almost surely. With Fatou's Lemma,

$$\mathbf{E}[|X|^p] = \mathbf{E}[\liminf_{k \rightarrow \infty} |X_{n_k}|^p] \leq \sup_{n \in \mathbb{N}} \mathbf{E}[|X_n|^p] < \infty$$

because of Lemma 7.9. In particular, $X \in \mathcal{L}^p$. Just like in Example 7.10, $\{|X_n - X|^p : n \in \mathbb{N}\}$ is uniformly integrable. For every $\delta > 0$, due to convergence in probability,

$$\mathbf{P}(|X_n - X| > \delta) \xrightarrow{n \rightarrow \infty} 0.$$

From lemma 7.9 now follows with dominated convergence

$$\lim_{n \rightarrow \infty} \mathbf{E}[|X_n - X|^p] = \lim_{n \rightarrow \infty} \mathbf{E}[|X_n - X|^p; |X_n - X| > \delta] + \mathbf{E}[|X_n - X|^p; |X_n - X| \leq \delta] \leq \delta^p.$$

Since $\delta > 0$ was arbitrary, $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$ follows. \square

Corollary 7.12 (\mathcal{L}^p -convergence and uniform integrability). *Let $1 \leq p < \infty$ and $X_1, X_2, \dots \in \mathcal{L}^p$ and X be measurable with $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$. Then, the following are equivalent:*

1. $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^p X$,
2. $\|X_n\|_p \xrightarrow{n \rightarrow \infty} \|X\|_p$,
3. The family $(|X_n|^p)_{n=1,2,\dots}$ is uniformly integrable.

Proof. The equivalence 1. \Leftrightarrow 3. is clear from Theorem 7.11.

1. \Rightarrow 2.: follows from Minkowski's inequality with

$$\left| \|X_n\|_p - \|X\|_p \right| \leq \|X_n - X\|_p \xrightarrow{n \rightarrow \infty} 0.$$

2. \rightarrow 3.: For fixed K , we write

$$\mathbf{E}[|X_n|^p; |X_n| > K] \leq \mathbf{E}[|X_n|^p - (|X_n| \wedge (K - |X_n|)^+)^p] \xrightarrow{n \rightarrow \infty} \mathbf{E}[|X|^p - (|X| \wedge (K - |X|)^+)^p].$$

Convergence follows from $\mathbf{E}[|X_n|^p] \xrightarrow{n \rightarrow \infty} \mathbf{E}[|X|^p]$, and $(|X_n| \wedge (K - |X_n|)^+)^p \xrightarrow{n \rightarrow \infty} \mathcal{L}^1 |X| \wedge (K - |X|)^+)^p$, since the convergence according to Proposition 7.6 is in probability, and $((|X_n| \wedge (K - |X_n|)^+)^p)_{n=1,2,\dots}$ is bounded, in particular uniformly integrable. Since $\mathbf{E}[|X|^p - (|X| \wedge (K - |X|)^+)^p] \xrightarrow{K \rightarrow \infty} 0$ after dominated convergence, $(|X_n|^p)_{n=1,2,\dots}$ is uniformly integrable. \square

8 Independence and the strong law

With our knowledge on probability measures and σ -algebras we now shed light on the concept of independence. In particular, in this chapter we will prove the strong law of large numbers, see Theorem 8.21. On the way, we prove the Borel-Cantelli lemma (Theorem 8.8) and Kolmogorov's 0-1 law (Theorem 8.15).

8.1 Definition and simple properties

Already in the lecture *Basic Probability*, independent random variables were considered. The intuitive idea of independence is often correct, but should sometimes be treated with caution.

Definition 8.1 (Independence). *1. A family of sets $(A_i)_{i \in I}$ with $A_i \in \mathcal{F}$ is called independent if*

$$\mathbf{P}\left(\bigcap_{j \in J} A_j\right) = \prod_{j \in J} \mathbf{P}(A_j) \tag{8.1}$$

for all $J \subseteq_f I$.²

²Recall that we write $J \subseteq_f I$ iff $J \subseteq I$ and J is finite.

2. A family $(\mathcal{C}_i)_{i \in I}$ of set systems $\mathcal{C}_i \subseteq \mathcal{F}$ is called independent if (8.1) holds for all $J \subseteq_f I$ and $A_j \in \mathcal{C}_j, j \in J$.

3. A family of random variables $(X_i)_{i \in I}$ is called independent if $(\sigma(X_i))_{i \in I}$ is independent.

We first deal with the question if there are probability spaces with an arbitrary number of independent random variables. Here we benefit from our knowledge of product measures.

Proposition 8.2 (Independence and product measures). *A family $(X_i)_{i \in I}$ of random variables is independent iff for each $J \subseteq_f I$*

$$((X_i)_{i \in J})_* \mathbf{P} = \bigotimes_{i \in J} (X_i)_* \mathbf{P},$$

i.e. the joint distribution of each finite subfamily is the product distribution of the individual distributions.

Proof. By definition, the family $(X_i)_{i \in I}$ is independent if and only if for each $J \subseteq_f I$ and $A_i \in \mathcal{F}, i \in J$,

$$\mathbf{P}(X_i \in A_i, i \in J) = \prod_{i \in J} \mathbf{P}(X_i \in A_i).$$

The assertion now follows from the fact that $\mathbf{P}(X_i \in A_i) = (X_i)_* \mathbf{P}(A_i)$ (see Definition ??) and $\mathbf{P}(X_i \in A_i, i \in J) = ((X_i)_{i \in J})_* \mathbf{P}(\times_{i \in J} A_i)$ (see Corollary 5.14). \square

Corollary 8.3 (Existence of uncountably many independent random variables). *Let E be a Polish space and I an arbitrary index set. Let $(\Omega_i, \mathcal{F}_i, \mathbf{P}_i)$ be probability spaces and X_i an E -valued random variable, $i \in I$. Then there is a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ and a family $(Y_i)_{i \in I}$ E -valued, independent random variable with $Y_i \stackrel{d}{=} X_i$.*

Proof. It should be noted that $((X_i)_{i \in J})_* \bigotimes_{i \in J} \mathbf{P}_i)_{J \subseteq_f I}$ is a projective family of probability measures on $(E, \mathcal{B}(E))$. Using Theorem 5.24 we find the projective limit \mathbf{P}_I . This is a probability measure on $(E^I, (\mathcal{B}(E))^I)$. Furthermore, with $\pi_i : E^I \rightarrow E$, the i -th projection, $(\pi_i)_* \mathbf{P}_I = (X_i)_* \mathbf{P}_i$, i.e. $\pi_i \stackrel{d}{=} X_i$. \square

Lemma 8.4 (Functions of independent random variables). *Let $(\Omega'_i, \mathcal{F}'_i), (\Omega''_i, \mathcal{F}''_i)$, $i \in I$, measurable spaces. Let $(X_i)_{i \in I}$ be a family of independent random variables, $X_i : \Omega \rightarrow \Omega'_i$, and $\varphi_i : \Omega'_i \rightarrow \Omega''_i$ measurable, $i \in I$. Then the family $(\varphi_i(X_i))_{i \in I}$ is independent.*

Proof. According to Lemma ??, the random variable $\varphi_i(X_i)$ is measurable according to $\sigma(X_i)$, $i \in I$, i.e. $\sigma(\varphi_i(X_i)) \subseteq \sigma(X_i)$. Since $(\sigma(X_i))_{i \in I}$ is an independent family by assumption, the assertion follows from the definition of independence. \square

Proposition 8.5 (Independent and Uncorrelated). *Let $X, Y \in \mathcal{L}^1$ be independent, real-valued random variables. Then $XY \in \mathcal{L}^1$ and*

$$\mathbf{E}[XY] = \mathbf{E}[X] \cdot \mathbf{E}[Y].$$

Proof. The assertion is clear if X and Y are indicator functions. Then, note that if the assertion applies to the pairs (X_i, Y_j) , $i, j = 1, \dots, n$, then it also for $\sum_{i=1}^n X_i$ and $\sum_{j=1}^n Y_j$: Indeed, due to the linearity of the expected value,

$$\mathbf{E}\left[\sum_{i=1}^n X_i \cdot \sum_{j=1}^n Y_j\right] = \sum_{i=1}^n \sum_{j=1}^n \mathbf{E}[X_i Y_j] = \sum_{i=1}^n \sum_{j=1}^n \mathbf{E}[X_i] \mathbf{E}[Y_j] = \mathbf{E}\left[\sum_{i=1}^n X_i\right] \cdot \mathbf{E}\left[\sum_{j=1}^n Y_j\right].$$

So, since the assertion applies to indicator functions, it is also valid for simple functions, and thus with monotonic convergence also for non-negative measurable functions. The general case follows with the decomposition $X = X^+ - X^-$ and $Y = Y^+ - Y^-$. \square

Example 8.6 (Uncorrelated, non-independent random variables). *Let U be a random variable uniformly distributed on $[0, 1]$, $X = \cos(2\pi U)$ and $Y = \sin(2\pi U)$. Then $\mathbf{E}[X] = \mathbf{E}[Y] = 0$ and*

$$\mathbf{E}[XY] = \int_0^1 \cos(2\pi u) \sin(2\pi u) du = \frac{1}{2} \int_0^1 \sin(4\pi u) du = 0$$

and thus X, Y are uncorrelated. However, $\{|X| < \varepsilon, |Y| < \varepsilon\} = \emptyset$ for $\varepsilon > 0$ is small enough and thus $\mathbf{P}(X^{-1}(-\varepsilon, \varepsilon), Y^{-1}(-\varepsilon, \varepsilon)) = 0 < \mathbf{P}(X^{-1}(-\varepsilon, \varepsilon)) \cdot \mathbf{P}(Y^{-1}(-\varepsilon, \varepsilon))$. This means that X and Y are not independent.

If there is a probability space and (countably) many events, you can ask yourself how many of these events will likely occur. The Borel-Cantelli lemma gives a sharp criterion for the occurrence of only finitely many events.

Definition 8.7 (Limsup of sets). *For $A_1, A_2, \dots \in \mathcal{F}$,*

$$\limsup_{n \rightarrow \infty} A_n := \bigcap_{n \geq 1} \bigcup_{m \geq n} A_m$$

is the event infinitely many of the A_n occur.

Theorem 8.8 (Borel-Cantelli lemma). *1. Let $A_1, A_2, \dots \in \mathcal{F}$. Then,*

$$\sum_{n=1}^{\infty} \mathbf{P}(A_n) < \infty \implies \mathbf{P}(\limsup_{n \rightarrow \infty} A_n) = 0.$$

2. If A_1, A_2, \dots are independent,

$$\sum_{n=1}^{\infty} \mathbf{P}(A_n) = \infty \implies \mathbf{P}(\limsup_{n \rightarrow \infty} A_n) = 1.$$

Proof. We start with 1. Because of the continuity of \mathbf{P} from above (see Proposition ??),

$$\mathbf{P}(\limsup_{n \rightarrow \infty} A_n) = \lim_{n \rightarrow \infty} \mathbf{P}\left(\bigcup_{m \geq n} A_m\right) \leq \lim_{n \rightarrow \infty} \sum_{m=n}^{\infty} \mathbf{P}(A_m) = 0$$

by assumption. For 2. we use that $\log(1-x) \leq -x$ for $x \in [0, 1]$. From this and the continuity of \mathbf{P} from below and the independence of $(A_n)_{n=1,2,\dots}$,

$$\begin{aligned}
\mathbf{P}((\limsup_{n \rightarrow \infty} A_n)^c) &= \mathbf{P}\left(\bigcup_{n=1}^{\infty} \bigcap_{m \geq n} A_m^c\right) \\
&= \lim_{n \rightarrow \infty} \mathbf{P}\left(\bigcap_{m=n}^{\infty} A_m^c\right) \\
&= \lim_{n \rightarrow \infty} \prod_{m=n}^{\infty} (1 - \mathbf{P}(A_m)) \\
&= \lim_{n \rightarrow \infty} \exp\left(\sum_{m=n}^{\infty} \log(1 - \mathbf{P}(A_m))\right) \\
&\leq \lim_{n \rightarrow \infty} \exp\left(-\sum_{m=n}^{\infty} \mathbf{P}(A_m)\right) \\
&= 0,
\end{aligned}$$

and the assertion follows. □

Example 8.9 (Infinite coin toss and geometric distributions).

1. We consider an infinite coin toss. This means that we have a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ and independent random variables X_1, X_2, \dots with values in $\{\text{heads}, \text{tails}\}$. The coin toss is fair, i.e. $\mathbf{P}(X_n = \text{head}) = 1/2$. We consider the events $A_n = \{X_n = \text{head}\}$. Since

$$\sum_{n=1}^{\infty} \mathbf{P}(A_n) = \sum_{n=1}^{\infty} \frac{1}{2} = \infty$$

and the family $(A_n)_{n \in \mathbb{N}}$ is independent, it follows from the Borel-Cantelli lemma that almost surely infinitely often head occurs.

2. We consider the same situation as in 1, but the events $B_n := \{X_1 = \text{Kopf}\}$. It is clear that the family $(B_n)_{n \in \mathbb{N}}$ is not independent. (For example $\mathbf{P}(B_1 \cap B_2) = \mathbf{P}(B_1) = 1/2 \neq \frac{1}{4} = \mathbf{P}(B_1) \cdot \mathbf{P}(B_2)$.) Just like in 1. $\sum_{n=1}^{\infty} \mathbf{P}(B_n) = \infty$. It is also clear that $\mathbf{P}(\limsup_{n \rightarrow \infty} B_n) = \frac{1}{2}$. It follows from this, that in the Borel-Cantelli lemma the condition of independence in 2. does not apply.
3. Let X_1, X_2, \dots be geometrically distributed with the success parameter p . We consider the events $A_n := \{X_n \geq n\}$ and ask ourselves whether an infinite number of these events can occur. Since

$$\sum_{n=1}^{\infty} \mathbf{P}(A_n) = \sum_{n=1}^{\infty} \mathbf{P}(X_n \geq n) = \sum_{n=1}^{\infty} (1-p)^{n-1} = \frac{1}{p} < \infty.$$

Therefore, almost certainly only a finite number of the events $\{X_n \geq n\}$ occur.

8.2 Kolmogorov's 0-1 law

The Borel-Cantelli lemma is already a statement about when an event that depends on an infinite number of events is almost certain to occur. We will now examine this situation further.

Proposition 8.10 (Independence of generated σ -algebras). *Let $(\mathcal{C}_i)_{i \in I}$ be a family of independent, \cap -stable set systems. Then, $(\sigma(\mathcal{C}_i))_{i \in I}$ is also an independent family.*

Proof. Let $J = \{i_1, \dots, i_n\} \subseteq_f I$ and (wlog) $n > 1$. Then, (8.1) holds for any A_{i_1}, \dots, A_{i_n} with $A_{i_k} \in \mathcal{C}_{i_k}, k = 1, \dots, n$. We keep A_{i_2}, \dots, A_{i_n} fixed and define

$$\mathcal{D} := \{A_{i_1} \in \mathcal{F} : (8.1) \text{ holds}\}.$$

We will now show that \mathcal{D} is a Dynkin system. Namely, if $A \subseteq B \in \mathcal{D}$, then $B \setminus A \in \mathcal{D}$, because

$$\begin{aligned} \mathbf{P}\left((B \setminus A) \cap \bigcap_{k=2}^n A_{i_k}\right) &= \mathbf{P}\left(B \cap \bigcap_{k=2}^n A_{i_k}\right) - \mathbf{P}\left(A \cap \bigcap_{k=2}^n A_{i_k}\right) \\ &= (\mathbf{P}(B) - \mathbf{P}(A)) \cdot \prod_{k=2}^n \mathbf{P}(A_{i_k}) \\ &= \mathbf{P}(B \setminus A) \cdot \prod_{k=2}^n \mathbf{P}(A_{i_k}). \end{aligned}$$

Furthermore, if $A_1, A_2, \dots \in \mathcal{D}$ with $A_1 \subseteq A_2 \subseteq A_3 \dots$, then due to the continuity of \mathbf{P} from below,

$$\begin{aligned} \mathbf{P}\left(\left(\bigcup_{j=1}^{\infty} A_j\right) \cap \bigcap_{k=2}^n A_{i_k}\right) &= \sup_{j \in \mathbb{N}} \mathbf{P}\left(A_j \cap \bigcap_{k=2}^n A_{i_k}\right) \\ &= \sup_{j \in \mathbb{N}} \mathbf{P}(A_j) \cdot \prod_{k=2}^n \mathbf{P}(A_{i_k}) \\ &= \mathbf{P}\left(\bigcup_{j=1}^{\infty} A_j\right) \cdot \prod_{k=2}^n \mathbf{P}(A_{i_k}). \end{aligned}$$

Since \mathcal{C}_{i_1} is \cap -stable and $\mathcal{C}_{i_1} \subseteq \mathcal{D}$, $\sigma(\mathcal{C}_{i_1}) \subseteq \mathcal{D}$ according to theorem 1.13. In particular, (8.1) applies for $A_{i_1} \in \sigma(\mathcal{C}_{i_1}), A_{i_2} \in \mathcal{C}_{i_2}, \dots, A_{i_n} \in \mathcal{C}_{i_n}$. Iterating the above procedure for $k = 2, \dots, n$, you get the statement. \square

Corollary 8.11 (Independence of indicator functions). *A family of sets $(A_i)_{i \in I}$ is independent if and only if the family of random variables $(1_{A_i})_{i \in I}$ is independent. In particular,*

$$\mathbf{P}\left(\bigcap_{j \in J} B_j\right) = \prod_{j \in J} \mathbf{P}(B_j)$$

for $J \subseteq_f I, B_j \in \{A_j, A_j^c\}, j \in J$.

Proof. For $i \in I$ let $\mathcal{C}_i = \{A_i\}$. Then $\sigma(1_{A_i}) = \{\emptyset, A_i, A_i^c, \Omega\} = \sigma(\mathcal{C}_i)$. Since \mathcal{C}_i is trivially cut-stable, the statement follows from Proposition 8.10. \square

Corollary 8.12 (Grouping). *Let $(\mathcal{F}_i)_{i \in I}$ be a family of independent σ -algebras. Further, let \mathcal{I} be a partition of I , i.e. $\mathcal{I} = \{I_k, k \in K\}$ with $\bigsqcup_{k \in K} I_k = I$, so the I_k are disjoint and their union is I . Then, $(\sigma(\mathcal{F}_i : i \in I_k))_{k \in K}$ is also an independent system.*

Proof. The set system $\mathcal{C}_k := \{\bigcap_{i \in J_k} A_i : J_k \subseteq_f I_k, A_i \in \mathcal{F}_i\}$ is \cap -stable and $\sigma(\mathcal{C}_k) = \sigma(\mathcal{F}_i : i \in I_k)$, $k \in K$. Since, according to the assumption, the family $(\mathcal{C}_k)_{k \in K}$ is independent, the assertion follows from Proposition 8.10. \square

We now come to the main statement of this section, Kolmogorov's 0-1 law. For this we introduce a certain σ -algebra, the terminal σ -algebra.

Definition 8.13 (Terminal and trivial σ -algebras). *1. Let $\mathcal{F}_1, \mathcal{F}_2, \dots \subseteq \mathcal{F}$ be a sequence of σ -algebras. Then*

$$\mathcal{T}(\mathcal{F}_1, \mathcal{F}_2, \dots) = \bigcap_{n \geq 1} \sigma\left(\bigcup_{m > n} \mathcal{F}_m\right)$$

the σ -algebra of terminal events of $\mathcal{F}_1, \mathcal{F}_2, \dots$

*2. A σ -algebra $\tilde{\mathcal{F}} \subseteq \mathcal{F}$ is called **P**-trivial if $\mathbf{P}(A) \in \{0, 1\}$ for all $A \in \tilde{\mathcal{F}}$.*

Lemma 8.14 (Trivial σ -algebras). *1. A σ -algebra $\tilde{\mathcal{F}}$ is **P**-trivial if and only if $\tilde{\mathcal{F}}$ is independent of itself.*

*2. Let $\tilde{\mathcal{F}}$ be a **P**-trivial σ -algebra and X a $\tilde{\mathcal{F}}$ -measurable random variable with values in a separable metric space E . Then X is constant, almost surely.*

Proof. 1. Let $\tilde{\mathcal{F}}$ be **P**-trivial and $A, B \in \tilde{\mathcal{F}}$. Then $\mathbf{P}(A \cap B) = \mathbf{P}(A) \wedge \mathbf{P}(B) = \mathbf{P}(A) \cdot \mathbf{P}(B)$, therefore $\tilde{\mathcal{F}}$ is independent of itself. If on the other hand, $\tilde{\mathcal{F}}$ is independent of itself and $A \in \tilde{\mathcal{F}}$, then $\mathbf{P}(A) = \mathbf{P}(A \cap A) = (\mathbf{P}(A))^2$, i.e. $\mathbf{P}(A) \in \{0, 1\}$.

2. For $n \in \mathbb{N}$, let $(B_{nj})_{j=1,2,\dots}$ be a countable covering of E with balls of radius $1/n$. Since $\tilde{\mathcal{F}}$ is a **P**-trivial σ -algebra then $\mathbf{P}(X \in B_{nj}) \in \{0, 1\}$ applies to all n, j . For $n \in \mathbb{N}$ let $J_n := \{j \in \mathbb{N} : \mathbf{P}(X \in B_{nj}) = 1\} \neq \emptyset$. Thus, due to the continuity from above, $\mathbf{P}\left(X \in \bigcap_{n=1}^{\infty} \bigcap_{j \in J_n} B_{nj}\right) = 1$. Since $\bigcap_{n=1}^{\infty} \bigcap_{j \in J_n} B_{nj}$ has at most one element, the assertion follows. \square

Under independence, the σ -algebra of terminal events is particularly simple.

Theorem 8.15 (Kolmogorov's 0-1 law). *Let $\mathcal{F}_1, \mathcal{F}_2, \dots \subseteq \mathcal{F}$ be a sequence of independent σ -algebras. Then $\mathcal{T} := \mathcal{T}(\mathcal{F}_1, \mathcal{F}_2, \dots)$ **P**-trivial.*

Proof. Let $\mathcal{T}_n := \sigma\left(\bigcup_{m > n} \mathcal{F}_m\right)$, $n = 1, 2, \dots$. According to corollary ??, $(\mathcal{F}_1, \dots, \mathcal{F}_n, \mathcal{T}_n)$ are independent, $n = 1, 2, \dots$. This means that $(\mathcal{F}_1, \dots, \mathcal{F}_n, \mathcal{T})$ are also independent, $n = 1, 2, \dots$ and thus also $(\mathcal{T}, \mathcal{F}_1, \mathcal{F}_2, \dots)$. Again with Corollary 8.12, it follows that $(\mathcal{T}_0, \mathcal{T})$ are independent and, since $\mathcal{T} \subseteq \mathcal{T}_0$ it also follows that \mathcal{T} is independent of itself. Therefore, the assertion follows from Lemma 8.14. \square

8.3 Sums of independent random variables

Many important theorems in probability theory deal with independent random variables. In this lecture, these are in particular the Strong Law of Large Numbers (Theorem 8.21) and the Central Limit Theorem (Theorem 10.8). We present here important tools for analyzing sums of independent random variables. The first is the connection with the convolution of probability measures (see section 5.4).

Proposition 8.16 (Convolution is distribution of the independent sum). *Let X_1, \dots, X_n be independent, real-valued random variables. Then,*

$$(X_1 + \dots + X_n)_* \mathbf{P} = (X_1)_* \mathbf{P} * \dots * (X_n)_* \mathbf{P}.$$

Further, for the characteristic functions

$$\psi_{X_1 + \dots + X_n} = \psi_{X_1} \cdots \psi_{X_n}$$

and, if X_1, \dots, X_n assume values in \mathbb{R}_+ ,

$$\mathcal{L}_{X_1 + \dots + X_n} = \mathcal{L}_{X_1} \cdots \mathcal{L}_{X_n}.$$

Proof. First of all, according to Proposition 8.2 $((X_1, \dots, X_n))_* \mathbf{P} = (X_1)_* \mathbf{P} \otimes \dots \otimes (X_n)_* \mathbf{P}$. Thus, the first assertion already follows from Definition ?? of the convolution of measures. The further assertions follow from Proposition ??, since for example

$$\begin{aligned} \psi_{X_1 + \dots + X_n}(t) &= \mathbf{E}[e^{it(X_1 + \dots + X_n)}] = \mathbf{E}[e^{itX_1} \cdots e^{itX_n}] \\ &= \mathbf{E}[e^{itX_1}] \cdots \mathbf{E}[e^{itX_n}] = \psi_{X_1}(t) \cdots \psi_{X_n}(t). \end{aligned} \quad \square$$

Kolmogorov's 0-1 law provides a very simple statement as to when sums of independent random variables are almost certain to converge.

Proposition 8.17 (Convergence of sums of independent random variables). *Let X_1, X_2, \dots be independent random variables and $S_n := X_1 + \dots + X_n$.*

1. *Then,*

$$\mathbf{P}(\omega : S_n(\omega) \text{ converges for } n \rightarrow \infty) \in \{0, 1\}$$

2. *Further,*

$$\mathbf{P}(\omega : S_n(\omega)/n \text{ converges for } n \rightarrow \infty) \in \{0, 1\}.$$

If $\mathbf{P}(S_n/n \text{ converges}) = 1$, the limit value is almost certainly constant.

Proof. Set $\mathcal{F}_i := \sigma(X_i)$, $i = 1, 2, \dots$. This means that the family $(\mathcal{F}_i)_{i=1,2,\dots}$ is independent. The set $\{\omega : S_n(\omega) \text{ converges for } n \rightarrow \infty\}$ is measurable with respect to $\mathcal{T}(\mathcal{F}_1, \mathcal{F}_2, \dots)$ and thus the first statement from Theorem 8.15 follows. In the same way it follows that $\mathbf{P}(S_n/n \text{ converges}) \in \{0, 1\}$. Let $S = \lim_{n \rightarrow \infty} S_n(n)/n$. Thus, for all $m = 1, 2, \dots$,

$$S = \lim_{n \rightarrow \infty} \frac{X_1 + \dots + X_n}{n} = \lim_{n \rightarrow \infty} \frac{X_m + \dots + X_n}{n},$$

so S is measurable wrt $\sigma\left(\bigcup_{k \geq m} \mathcal{F}_k\right)$. This means that S is also \mathcal{T} -measurable and therefore almost certainly constant according to Theorem 8.15 and Lemma 8.14. \square

Proposition 8.18 (Maximum inequality of Kolmogorov). *Let $X_1, X_2, \dots \in \mathcal{L}^2$ be independent random variables. Then, for $K > 0$,*

$$\mathbf{P}\left(\sup_{n \in \mathbb{N}} \left| \sum_{k=1}^n X_k - \mathbf{E}[X_k] \right| > K\right) \leq \frac{\sum_{n=1}^{\infty} \mathbf{V}(X_n)}{K^2}.$$

Proof. Wlog, let $\mathbf{E}[X_k] = 0, k = 1, 2, \dots$. We further set $S_n = X_1 + \dots + X_n$ and $T := \inf\{n : |S_n| > K\}$. Then, $\mathbf{P}(\sup_n |S_n| > K) = \mathbf{P}(T < \infty)$. Because of Corollary 8.12, $S_k \cdot 1_{T=k}$ and $S_n - S_k$ are independent for $k \leq n$. Therefore

$$\begin{aligned} \sum_{k=1}^n \mathbf{E}[X_k^2] &= \mathbf{E}[S_n^2] \geq \sum_{k=1}^n \mathbf{E}[S_n^2, T = k] \\ &= \sum_{k=1}^n \mathbf{E}[S_k^2 + (S_n - S_k + 2S_k)(S_n - S_k), T = k] \\ &\geq \sum_{k=1}^n \mathbf{E}[S_k^2, T = k] + 2\mathbf{E}[S_k(S_n - S_k), T = k] \\ &= \sum_{k=1}^n \mathbf{E}[S_k^2, T = k] \geq K^2 \mathbf{P}(T \leq n) \end{aligned}$$

Now follows the assertion with $n \rightarrow \infty$. □

Theorem 8.19 (Convergence criterion for series). *Let $X_1, X_2, \dots \in \mathcal{L}^2$ be independent random variables with $\sum_{n=1}^{\infty} \mathbf{V}[X_n] < \infty$. Then, $\sum_{k=1}^n X_k - \mathbf{E}[X_k]$ converges almost surely.*

Proof. Again, let $\mathbf{E}[X_k] = 0, k = 1, 2, \dots$ and we write $S_n = X_1 + \dots + X_n$. For $\varepsilon > 0$, according to Proposition 8.18,

$$\lim_{k \rightarrow \infty} \mathbf{P}(\sup_{n \geq k} |S_n - S_k| > \varepsilon) \leq \lim_{k \rightarrow \infty} \frac{\sum_{n=k+1}^{\infty} \mathbf{E}[X_n^2]}{\varepsilon^2} = 0.$$

Therefore, $\sup_{n \geq k} |S_n - S_k| \xrightarrow{k \rightarrow \infty} 0$. So, by Proposition 7.6, there is a subsequence k_1, k_2, \dots with $\sup_{n \geq k_i} |S_n - S_{k_i}| \xrightarrow{i \rightarrow \infty} 0$. However, since $(\sup_{n \geq k} |S_n - S_k|)_{k=1,2,\dots}$ is decreasing, $\sup_{n \geq k} |S_n - S_k| \xrightarrow{k \rightarrow \infty} 0$ applies. This means, however, that $(S_n)_{n=1,2,\dots}$ converges. □

8.4 The Strong Law of Large Numbers

In the lecture *Basic Probability*, we already proved the weak law of large numbers: if $X_1, X_2, \dots \in \mathcal{L}^2$ are identically distributed and uncorrelated, then, for $\varepsilon > 0$

$$\mathbf{P}\left(\frac{1}{n} \left| \sum_{k=1}^n (X_k - \mathbf{E}[X_k]) \right| > \varepsilon\right) \leq \frac{1}{\varepsilon^2} \mathbf{V}\left[\frac{1}{n} \sum_{k=1}^n X_k\right] = \frac{1}{\varepsilon^2 n^2} \sum_{k=1}^n \mathbf{V}[X_k] = \frac{\mathbf{V}[X_1]}{\varepsilon^2 n} \xrightarrow{n \rightarrow \infty} 0.$$

As we now know, this means in other terms,

$$\frac{1}{n} \sum_{k=1}^n X_k \xrightarrow{n \rightarrow \infty} \mathbf{E}[X_0].$$

We now want to improve this statement in two directions. On the one hand, we want to replace convergence in probability by almost sure convergence, and on the other hand only assume the existence of first moments (but not the existence of second moments). First, however, we define what exactly what we mean when we say that a sequence of random variables follows a law of large numbers.

Definition 8.20 (Law of large numbers). *Let $X_1, X_2, \dots \in \mathcal{L}^1$ be a sequence of real-valued random variables. We say that the sequence follows the weak law of large numbers if*

$$\frac{1}{n} \sum_{k=1}^n (X_k - \mathbf{E}[X_k]) \xrightarrow[n \rightarrow \infty]{p} 0.$$

The sequence satisfies the strong law of large numbers if

$$\frac{1}{n} \sum_{k=1}^n (X_k - \mathbf{E}[X_k]) \xrightarrow[n \rightarrow \infty]{fs} 0.$$

Theorem 8.21 (Strong law for independent random variables). *A sequence $X_1, X_2, \dots \in \mathcal{L}^1$ of independent and identically distributed random variables satisfies the strong law of large numbers, i.e.*

$$\frac{1}{n} \sum_{k=1}^n X_k \xrightarrow[n \rightarrow \infty]{fs} \mathbf{E}[X_1].$$

Remark 8.22 (Weak law of large numbers). *Since convergence in probability is implied by almost sure convergence (see Proposition 7.6), the sequence X_1, X_2, \dots from the theorem also satisfies the weak law of large numbers. Furthermore, the sequence X_1^+, X_2^+, \dots also satisfies the strong law and $\mathbf{E}[\frac{1}{n}(X_1^+ + \dots + X_n^+)] = \mathbf{E}[X_1^+]$. This means that the sequence $(\frac{1}{n}(X_1^+ + \dots + X_n^+))_{n=1,2,\dots}$ is uniformly according to Corollary 7.12. In the same way, the sequence of partial sums of the negative parts is uniformly integrable. It follows from Theorem 7.11 that $\frac{1}{n}(X_1 + \dots + X_n) \xrightarrow[n \rightarrow \infty]{\mathcal{L}^1} \mathbf{E}[X_1]$.*

Remark 8.23 (Finite fourth and second moments). *The difficulty in proving the strong law is that only may be used that $X_1 \in \mathcal{L}^1$. The proof is significantly easier if we use $X_1 \in \mathcal{L}^4$ or $X_1 \in \mathcal{L}^2$. We start with these two proofs and write $S_n := X_1 + \dots + X_n$.*

1. *The case $X_1 \in \mathcal{L}^4$: Here you can get by without further aids: From the linearity of the expected value, it is clear that $\mathbf{E}[S_n/n] = \mathbf{E}[X_1]$. Wlog, let $\mathbf{E}[X_1] = 0$, otherwise you go to the random variables $X_1 - \mathbf{E}[X_1], X_2 - \mathbf{E}[X_2], \dots \in \mathcal{L}^4$. First we calculate with the help of the independence of $(X_k)_{k=1,2,\dots}$*

$$\mathbf{E}[S_n^4] = \sum_{k=1}^n \mathbf{E}[X_k^4] + 3 \sum_{\substack{k,l=1 \\ k \neq l}}^{\infty} \mathbf{E}[X_k^2 X_l^2] \leq (n + 6n^2) \mathbf{E}[X_1^4]$$

because of the Cauchy-Schwartz inequality. From this,

$$\mathbf{E} \left[\sum_{n=1}^{\infty} \left(\frac{S_n}{n} \right)^4 \right] \leq \sum_{n=1}^{\infty} \frac{n + 6n^2}{n^4} \mathbf{E}[X_1^4] < \infty.$$

Therefore, $\sum_{n=1}^{\infty} \left(\frac{S_n}{n} \right)^4 < \infty$ applies almost certain, in particular $\frac{S_n}{n} \xrightarrow[n \rightarrow \infty]{fs} 0$.

2. The case $X_1 \in \mathcal{L}^2$: Here the convergence criterion for series, theorem 8.19 is of crucial help. We also need the following result:

Lemma 8.24 (Kronecker Lemma). *Let $x_1, x_2, \dots \in \mathbb{R}$, $y_1, y_2, \dots \in \mathbb{R}$ be monotone with $y_n \uparrow \infty$ and $\sum_{n=1}^{\infty} x_n/y_n < \infty$. Then, $\sum_{k=1}^n x_k/y_n \xrightarrow{n \rightarrow \infty} 0$.*

Proof. Let $z_0 = 0, z_n := \sum_{k=1}^n x_k/y_k$. Then $z_n \xrightarrow{n \rightarrow \infty} z_\infty < \infty$ and $x_k = y_k(z_k - z_{k-1})$. We write with $y_0 = 0$

$$\begin{aligned} \frac{\sum_{k=1}^n x_k}{y_n} &= \frac{1}{y_n} \sum_{k=1}^n y_k(z_k - z_{k-1}) = z_n + \frac{1}{y_n} \left(\sum_{k=0}^{n-1} y_k z_k - \sum_{k=1}^n y_k z_{k-1} \right) \\ &= z_n - \frac{1}{y_n} \left(\sum_{k=1}^n y_k z_{k-1} - y_{k-1} z_{k-1} \right) \\ &\xrightarrow{n \rightarrow \infty} z_\infty - z_\infty \cdot \lim_{n \rightarrow \infty} \frac{1}{y_n} \sum_{k=1}^n y_k - y_{k-1} = 0. \end{aligned}$$

□

Back to the proof of the strong law in the case $X_1 \in \mathcal{L}^2$. Wlog, let $\mathbf{E}[X_1] = 0$. Consider the sequence $X_1/1, X_2/2, \dots$. Because $\sum_{n=1}^{\infty} \mathbf{V}[X_n/n] = \mathbf{V}[X_1] \sum_{n=1}^{\infty} 1/n^2$ applies according to Theorem 8.19 that $\sum_{k=1}^n X_k/k$ almost surely converges. With Lemma 8.24 it follows that $S_n/n \xrightarrow{n \rightarrow \infty}_{fs} 0$.

Proof of theorem 8.21 if $X_1 \in \mathcal{L}^1$. It is sufficient to consider the case of non-negative random variables. In the general case, note that $X_1^+, X_2^+, \dots \in \mathcal{L}^1$ and $X_1^-, X_2^-, \dots \in \mathcal{L}^1$ fulfill the conditions of the theorem, and from $(X_1^+ + \dots + X_n^+)/n \xrightarrow{n \rightarrow \infty}_{fs} \mathbf{E}[X_1^+]$ and $(X_1^- + \dots + X_n^-)/n \xrightarrow{n \rightarrow \infty}_{fs} \mathbf{E}[X_1^-]$ the statement follows due to linearity of the expectation.

For $S_n = X_1 + \dots + X_n$ we will show that

$$\mathbf{E}[\limsup_{n \rightarrow \infty} S_n/n] \leq \mathbf{E}[X_1]. \quad (8.2)$$

If this is true, then firstly

$$\begin{aligned} \mathbf{E}[\liminf_{n \rightarrow \infty} S_n/n] &\geq \mathbf{E}[\liminf_{n \rightarrow \infty} (X_1 \wedge k + \dots + X_n \wedge k)/n] \\ &= k - \mathbf{E}[\limsup_{n \rightarrow \infty} ((k - X_1)^+ + \dots + (k - X_n)^+)/n] \\ &\geq \mathbf{E}[k - (k - X_1)^+] \xrightarrow{k \rightarrow \infty} \mathbf{E}[X_1]. \end{aligned}$$

Secondly, then $\mathbf{E}[\limsup_{n \rightarrow \infty} S_n/n - \liminf_{n \rightarrow \infty} S_n/n] = 0$, i.e. $\limsup_{n \rightarrow \infty} S_n/n = \liminf_{n \rightarrow \infty} S_n/n = 0$ almost surely, since both $\liminf_{n \rightarrow \infty} S_n/n$ as well as $\limsup_{n \rightarrow \infty} S_n/n$ are terminal functions, and thus according to Theorem 8.15 and Lemma 8.14 are almost surely constant. Furthermore,

$$\liminf_{n \rightarrow \infty} S_n/n = \mathbf{E}[\liminf_{n \rightarrow \infty} S_n/n] \geq \mathbf{E}[X_1] \geq \mathbf{E}[\limsup_{n \rightarrow \infty} S_n/n] = \limsup_{n \rightarrow \infty} S_n/n,$$

from which the assertion follows.




Figure 1: Illustration of M_i, I_j^n , introduced below (8.3). The size L_n is the number of contiguous areas of \bullet 's.

It therefore remains to show (8.2). Wlog let $\mathbf{E}[X_1] > 0$, otherwise $X_k = 0$ is almost certain, $k = 1, 2, \dots$ and the statement is trivial. For this we will use

$$0 < \alpha < \mathbf{E}[\limsup_{n \rightarrow \infty} S_n/n] \implies \alpha \leq \mathbf{E}[X_1] \quad (8.3)$$

can be proved. According to the assumption, for $i = 0, 1, 2, \dots$

$$\alpha < \mathbf{E}[\limsup_{n \rightarrow \infty} S_n/n] = \limsup_{n \rightarrow \infty} S_n/n = \limsup_{n \rightarrow \infty} (X_{i+1} + \dots + X_{i+n})/n.$$

Thus,

$$M_i := \inf\{n \in \mathbb{N} : (X_i + \dots + X_{i+n-1})/n \geq \alpha\}$$

is finite, almost surely, $i = 1, 2, \dots$. The M_i 's are identically distributed. We define recursively for $n = 1, 2, \dots$ (see also Figure 1) $I_1^n = 0$ and for $j = 0, 1, 2, \dots$ (with $M_0 := 0$)

$$I_{j+1}^n := \inf\{i \in \mathbb{N} : i \geq I_j^n + M_{I_j^n}, M_i \leq n - i + 1\}$$

with $\inf \emptyset = \infty$ and $L_n := \sup\{n \in \mathbb{N}_0 : I_j^n < \infty\}$. This means that for $1 \leq j \leq L_n$, $I_j^n + M_{I_j^n} \leq n$, i.e. $(X_{I_j^n} + \dots + X_{I_j^n + M_{I_j^n} - 1})/M_{I_j^n} \geq \alpha$. We now use this by means of

$$\begin{aligned} \mathbf{E}[X_1] &= \mathbf{E}[(X_1 + \dots + X_n)/n] \\ &\geq \frac{1}{n} \mathbf{E} \left[\sum_{j=1}^{L_n} M_{I_j^n} \cdot (X_{I_j^n} + \dots + X_{I_j^n + M_{I_j^n} - 1}) / M_{I_j^n} \right] \\ &\geq \frac{\alpha}{n} \mathbf{E} \left[\sum_{j=1}^{L_n} M_{I_j^n} \right] = \alpha - \frac{\alpha}{n} \mathbf{E} \left[n - \sum_{j=1}^{L_n} M_{I_j^n} \right] \\ &\geq \alpha - \frac{\alpha}{n} \mathbf{E} \left[\sum_{i=1}^n 1_{M_i > n-i+1} \right] \\ &= \alpha \left(1 - \frac{1}{n} \sum_{i=1}^n \mathbf{P}(M_i > i) \right) \xrightarrow{n \rightarrow \infty} \alpha, \end{aligned}$$

since $(\frac{1}{n} \sum_{i=1}^n \mathbf{P}(M_i > i))_{n=1,2,\dots}$ as Cesàro-Limes of $(\mathbf{P}(M_i > i))_{i=1,2,\dots}$ because of the identity of the distributions of the M_i 's converges to 0. Thus (8.3) is shown and the assertion is proven. \square

We now give a simple application of the strong law. It often happens in statistics that a large number of independent, identically distributed, real-valued random variables must be studied. The Glivenko-Cantelli theorem (Theorem 8.26) states that the empirical distribution of the random variables almost surely converges to the underlying distribution.

Definition 8.25 (Empirical distribution). *Let X_1, X_2, \dots be random variables. For $n = 1, 2, \dots$ the distribution is called (random) probability distribution*

$$\widehat{\mu}_n := \frac{1}{n} \sum_{k=1}^n \delta_{X_k}$$

the empirical distribution of X_1, \dots, X_n . If the random variables are real-valued, then in addition

$$\widehat{F}_n(x) := \frac{1}{n} \sum_{k=1}^n 1_{X_k \leq x},$$

the empirical distribution function of X_1, \dots, X_n .

Theorem 8.26 (Glivenko-Cantelli Theorem). *Let X_1, X_2, \dots be independent, real-valued random variables with identical distribution with distribution function F . Then,*

$$\lim_{n \rightarrow \infty} \sup_{x \in \mathbb{R}} \widehat{F}_n(x) - F(x) \xrightarrow{f.s.} 0.$$

Proof. For $x \in \mathbb{R}$ and $n = 1, 2, \dots$ let $Y_n(x) := 1_{X_n \leq x}$ and $Z_n(x) := 1_{X_n < x}$. According to Theorem 8.21, for each $x \in \mathbb{R}$

$$\begin{aligned} \widehat{F}_n(x) &= \frac{1}{n} \sum_{k=1}^n Y_k(x) \xrightarrow{f.s.} \mathbf{E}[Y_1(x)] = \mathbf{P}(X_1 \leq x) = F(x), \\ \widehat{F}_n(x-) &= \frac{1}{n} \sum_{k=1}^n Z_k(x) \xrightarrow{f.s.} \mathbf{E}[Z_1(x)] = \mathbf{P}(X_1 < x) = F(x-). \end{aligned}$$

We must show that these limits hold uniformly for all $x \in \mathbb{R}$. For $N = 1, 2, \dots$ and $j = 0, \dots, N$ we set

$$x_j^N := \inf\{x \in \mathbb{R} : F(x) \geq j/N\}$$

and

$$R_n^N := \max_{j=1, \dots, N-1} (|\widehat{F}_n(x_j^N) - F(x_j^N)| + |\widehat{F}_n(x_j^N-) - F(x_j^N-)|).$$

For $N = 1, 2, \dots$, therefore, $R_n^N \xrightarrow{f.s.} 0$. Furthermore, for $x \in (x_{j-1}^N, x_j^N)$

$$\begin{aligned} \widehat{F}_n(x) &\leq \widehat{F}_n(x_j^N) \leq \widehat{F}_n(x_j^N) + R_n^N \leq F(x) + R_n^N + \frac{1}{N}, \\ \widehat{F}_n(x) &\geq \widehat{F}_n(x_{j-1}^N) \geq F(x_{j-1}^N) - R_n^N \geq F(x) - R_n^N - \frac{1}{N}, \end{aligned}$$

thus, for each $N = 1, 2, \dots$

$$\sup_{x \in \mathbb{R}} \widehat{F}_n(x) - F(x) \leq \frac{1}{N} + R_n^N \xrightarrow{f.s.} \frac{1}{N}.$$

Since the left-hand side does not depend on N , the assertion follows with $N \rightarrow \infty$. □

9 Weak convergence

For measurable spaces, we have often used the Borel σ -algebra, i.e. the σ -algebra that is generated by a topology. In this section we will often assume that the topological space is Polish, i.e. separable and metrizable by a complete metric; recall from Definition A.1 in the manuscript on measure theory. To save us some work, we will assume throughout that (E, r) is a metric space and sometimes we will assume that it is complete and separable.

For a measurable mapping $f : E \rightarrow \mathbb{R}$ and a measure μ on $\mathcal{B}(E)$ (the Borel's σ -algebra of E) we will use throughout this and the next chapter the notation

$$\mu[f] := \int f d\mu.$$

9.1 Definition and simple properties

So far, we have dealt with different types of convergence of random variables. The convergence in distribution of random variables is the same as the weak convergence of the distributions of random variables. For the motivation behind the following definitions, let us recall a fact: in a metric space (E, r) we have $x_n \xrightarrow{n \rightarrow \infty} x$ if and only if $f(x_n) \xrightarrow{n \rightarrow \infty} f(x)$ for all continuous functions on E (i.e. $f \in \mathcal{C}(E, \mathbb{R})$).

Definition 9.1 (Weak convergence and convergence in distribution).

1. We denote by $\mathcal{P}(E)$ the set of probability measures on $\mathcal{B}(E)$ and with $\mathcal{P}_{\leq 1}(E)$ the set of finite measures μ on $\mathcal{B}(E)$ with $\mu(E) \leq 1$. Further, $\mathcal{C}_b(E)$ is the set of the real-valued, bounded, continuous functions on E and $\mathcal{C}_c(E) \subseteq \mathcal{C}_b(E)$ is the set of the real-valued, bounded continuous functions on E with compact support.
2. A sequence $\mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(E)$ converges weakly to $\mathbf{P} \in \mathcal{P}(E)$, if

$$\mathbf{P}_n[f] \xrightarrow{n \rightarrow \infty} \mathbf{P}[f] \tag{9.1}$$

for all $f \in \mathcal{C}_b(E)$. We then write

$$\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}.$$

3. Let $\mu_1, \mu_2, \dots \in \mathcal{P}_{\leq 1}$ and μ be a measure on E . If (9.1) only applies to all $f \in \mathcal{C}_c(E)$, we say that μ_n converges vaguely to μ . We then write

$$\mu_n \xrightarrow[n \rightarrow \infty]{v} \mu.$$

4. Let X, X_1, X_2, \dots be random variables on probability spaces $(\Omega, \mathcal{A}, \mathbf{P}), (\Omega_1, \mathcal{A}_1, \mathbf{P}_1), (\Omega_2, \mathcal{A}_2, \mathbf{P}_2), \dots$ with values in E . Then, X_1, X_2, \dots converges in distribution to X if $(X_n)_* \mathbf{P}_n \xrightarrow{n \rightarrow \infty} X_* \mathbf{P}$. We then write

$$X_n \xrightarrow{n \rightarrow \infty} X.$$

Remark 9.2. 1. Note that for random variables X, X_1, X_2, \dots with values in E , we have $X_n \xrightarrow{n \rightarrow \infty} X$ if

$$\mathbf{P}[f(X_n)] \xrightarrow{n \rightarrow \infty} \mathbf{P}[f(X)]$$

for all $f \in C_b(E)$. Many of the following results can therefore be formulated in two ways: either by means of probability distributions, or by means of random variables. The connection here is always that the statement about the probability distributions is also a statement about the distributions of the random variables.

2. The weak limit of probability measures must again be a probability measure, since $1 \in C_b(E)$. The vague limit of probability measures does not necessarily have to be a probability measure, since $1 \notin C_c(E)$ if E is not compact; see also Example 9.3.1. After all, the vague limit is in $\mathcal{P}_{\leq 1}(E)$, as Lemma 9.12 shows.
3. We already know the almost sure convergence, the convergence in probability, and the convergence in \mathcal{L}^p of random variables X_1, X_2, \dots to X . The difference to convergence in distribution is that the latter does not require that the random variables are defined on the same probability space.
4. By Definition 9.1, the topology of weak convergence on $\mathcal{P}(E)$ is the weakest (i.e. the smallest) topology for which $\mathbf{P} \mapsto \mathbf{P}[f]$ for all $f \in C_b(E)$ is continuous.

Example 9.3. 1. Let $x, x_1, x_2, \dots \in \mathbb{R}$ with $x_n \xrightarrow{n \rightarrow \infty} x$ and $\mathbf{P} = \delta_x, \mathbf{P}_1 = \delta_{x_1}, \mathbf{P}_2 = \delta_{x_2}, \dots$. Then, $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$, since

$$\mathbf{P}_n[f] = f(x_n) \xrightarrow{n \rightarrow \infty} f(x) = \mathbf{P}[f]$$

for all $f \in C_b(\mathbb{R})$.

If the sequence x_1, x_2, \dots diverges, for example $x_n = n$, then $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} 0$ (this is the 0-measure on $\mathcal{B}(\mathbb{R})$), since

$$\mathbf{P}_n[f] = f(x_n) \xrightarrow{n \rightarrow \infty} 0 = 0[f]$$

for all $f \in C_c(\mathbb{R})$. However, weak convergence does not hold, since $\mathbf{P}_n[1] = 1 \neq 0 = 0[1]$.

2. Let X, X_1, X_2, \dots be identically distributed. Then $X_n \xrightarrow{n \rightarrow \infty} X$, but in general the convergence is neither almost sure, nor in probability nor in \mathcal{L}^p for any $p > 0$.
3. As we will see, the Central Limit Theorem (Theorem 10.8), is a result about convergence in distribution. In its simplest form, the theorem of deMoivre-Laplace (see also Remark 9.8 and Example 9.34), it states: let $p \in (0, 1)$, $X_n \sim B(n, p), n = 1, 2, \dots$ and $X \sim N(0, 1)$. Then,

$$\frac{X_n - np}{\sqrt{np(1-p)}} \xrightarrow{n \rightarrow \infty} X.$$

4. Similarly, the Poisson approximation of the binomial distribution is a statement about convergence in distribution (see the course in Basic Probability and Theorem 10.5): let $X_n \sim B(n, p_n), n = 1, 2, \dots$ with $n \cdot p_n \xrightarrow{n \rightarrow \infty} \lambda$ and $X \sim Poi(\lambda)$. Then,

$$X_n \xrightarrow{n \rightarrow \infty} X.$$

Lemma 9.4 (Uniqueness of the weak limit). *Let $\mathbf{P}, \mathbf{Q}, \mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(E)$ with $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$ and $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{Q}$. Then $\mathbf{P} = \mathbf{Q}$.*

Proof. According to Proposition 2.11 it suffices to show that $\mathbf{P}(A) = \mathbf{Q}(A)$ for all closed $A \subseteq E$. (The set of all closed sets is a \cap -stable generator of $\mathcal{B}(E)$.) So let $A \subseteq E$ be closed. We set

$$r(x, A) := \inf_{y \in A} r(x, y)$$

and

$$f_m(x) \mapsto (1 - m \cdot r(x, A))^+.$$

for $m = 1, 2, \dots$. Then $f_m \xrightarrow{m \rightarrow \infty} 1_A$, since A is closed. Then, using dominated convergence,

$$\mathbf{P}(A) = \lim_{m \rightarrow \infty} \mathbf{P}[f_m] = \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbf{P}_n[f_m] = \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbf{Q}[f_m] = \mathbf{Q}(A)$$

and the assertion follows. \square

Recall the initial figure of Chapter 7. A sequence of random variables can converge almost surely, in probability, in \mathcal{L}^p or in distribution. Convergence in distribution is the weakest of these terms in the following sense.

Proposition 9.5 (Convergence in probability and in distribution). *Let X, X_1, X_2, \dots be random variables with values in E . If $X_n \xrightarrow{n \rightarrow \infty}_p X$, then $X_n \xrightarrow{n \rightarrow \infty} X$. If X is constant, the inversion also applies.*

Proof. Let $X_n \xrightarrow{n \rightarrow \infty}_p X$. Suppose that there is an $f \in \mathcal{C}_b(E)$ such that $\lim_{n \rightarrow \infty} \mathbf{P}[f(X_n)] \neq \mathbf{P}[f(X)]$. Then there is a subsequence $(n_k)_{k=1,2,\dots}$ and a $\varepsilon > 0$ with

$$\lim_{k \rightarrow \infty} |\mathbf{P}[f(X_{n_k})] - \mathbf{P}[f(X)]| > \varepsilon. \quad (9.2)$$

Because of $X_n \xrightarrow{n \rightarrow \infty}_p X$ and Proposition 7.6 there is a subsequence $(n_{k_\ell})_{\ell=1,2,\dots}$ such that $X_{n_{k_\ell}} \xrightarrow{\ell \rightarrow \infty} X$ almost surely. By dominated convergence, this would imply

$$\lim_{\ell \rightarrow \infty} \mathbf{P}[f(X_{n_{k_\ell}})] = \mathbf{P}[f(X)]$$

in contradiction to (9.2).

For the inverse, let $X = s \in E$. Note that $x \mapsto r(x, s) \wedge 1$ is a bounded, continuous function and therefore

$$\mathbf{P}[r(X_n, s) \wedge 1] \xrightarrow{n \rightarrow \infty} \mathbf{P}[r(X, s) \wedge 1] = 0.$$

Thus, $X_n \xrightarrow{n \rightarrow \infty}_p X$ holds because of (7.1). \square

Theorem 9.6 (Portmanteau theorem). *Let X, X_1, X_2, \dots be random variables with values in E . The following conditions are equivalent:*

(i) $X_n \xrightarrow{n \rightarrow \infty} X$

(ii) $\mathbf{P}[f(X_n)] \xrightarrow{n \rightarrow \infty} \mathbf{P}[f(X)]$ for all bounded, Lipschitz-continuous functions f .

(iii) $\liminf_{n \rightarrow \infty} \mathbf{P}(X_n \in G) \geq \mathbf{P}(X \in G)$ for all open $G \subseteq E$.

(iv) $\limsup_{n \rightarrow \infty} \mathbf{P}(X_n \in F) \leq \mathbf{P}(X \in F)$ for all completed $F \subseteq E$.

(v) $\lim_{n \rightarrow \infty} \mathbf{P}(X_n \in B) = \mathbf{P}(X \in B)$ for all $B \in \mathcal{B}(E)$ with³ $\mathbf{P}(X \in \partial B) = 0$.

Proof. (i) \rightarrow (ii): clear.

(ii) \Rightarrow (iv): Let $F \subseteq E$ be closed and f_1, f_2, \dots Lipschitz-continuous such that $f_k \downarrow 1_F$. (For example, one chooses $\varepsilon_k \downarrow 0$ and $f_k(x) = (1 - \frac{1}{\varepsilon_k} r(x, F))^+$, where $r(x, F) := \inf_{y \in F} r(x, y)$.) This means that

$$\limsup_{n \rightarrow \infty} \mathbf{P}(X_n \in F) \leq \inf_{k=1,2,\dots} \limsup_{n \rightarrow \infty} \mathbf{P}[f_k(X_n)] = \inf_{k=1,2,\dots} \mathbf{P}[f_k(X)] = \mathbf{P}(X \in F).$$

(iii) \iff (iv): That is clear. For (iii) \Rightarrow (iv), set $F := E \setminus G$ and for (iv) \Rightarrow (iii), set $G := E \setminus F$.

(iii) \Rightarrow (i): Let $f \geq 0$ be continuous. By Proposition 6.10 and Fatou's lemma,

$$\begin{aligned} \mathbf{P}[f(X)] &= \int_0^\infty \mathbf{P}(f(X) > t) dt \leq \int_0^\infty \liminf_{n \rightarrow \infty} \mathbf{P}(f(X_n) > t) dt \\ &\leq \liminf_{n \rightarrow \infty} \int_0^\infty \mathbf{P}(f(X_n) > t) dt = \liminf_{n \rightarrow \infty} \mathbf{P}[f(X_n)]. \end{aligned}$$

For $-c < f < c$, since $-f + c \geq 0$,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathbf{P}[f(X_n)] &= c - \liminf_{n \rightarrow \infty} \mathbf{P}[-f(X_n) + c] \leq c - \mathbf{P}[-f(X) + c] = \mathbf{P}[f(X)] \\ &\leq \liminf_{n \rightarrow \infty} \mathbf{P}[f(X_n)], \end{aligned}$$

thus $\mathbf{P}[f(X_n)] \xrightarrow{n \rightarrow \infty} \mathbf{P}[f(X)]$.

(iii), (iv) \rightarrow (v) For $B \in \mathcal{B}(E)$,

$$\mathbf{P}(X \in B^\circ) \leq \liminf_{n \rightarrow \infty} \mathbf{P}(X_n \in B^\circ) \leq \limsup_{n \rightarrow \infty} \mathbf{P}(X_n \in \overline{B}) \leq \mathbf{P}(X \in \overline{B}).$$

Given $\mathbf{P}(X \in \partial B) = \mathbf{P}(X \in \overline{B}) - \mathbf{P}(X \in B^\circ) = 0$, therefore $\mathbf{P}(X_n \in B) \xrightarrow{n \rightarrow \infty} \mathbf{P}(X \in B)$.

(v) \rightarrow (iv): Assume (v) is true and $F \subseteq E$ is closed. We write $F^\varepsilon := \{x \in E : r(x, F) \leq \varepsilon\}$ for $\varepsilon > 0$. The sets $\partial F^\varepsilon \subseteq \{x : r(x, F) = \varepsilon\}$ are disjoint, so

$$\mathbf{P}(X \in \partial F^\varepsilon) = 0 \tag{9.3}$$

for Lebesgue-almost every ε . Let $\varepsilon_1, \varepsilon_2, \dots$ denote a sequence with $\varepsilon_k \downarrow 0$ such that (9.3) holds for all $\varepsilon_1, \varepsilon_2, \dots$. This means that

$$\limsup_{n \rightarrow \infty} \mathbf{P}(X_n \in F) \leq \inf_{k=1,2,\dots} \limsup_{n \rightarrow \infty} \mathbf{P}(X_n \in F^{\varepsilon_k}) = \inf_{k=1,2,\dots} \mathbf{P}(X \in F^{\varepsilon_k}) = \mathbf{P}(X \in F).$$

□

³For the closure \overline{B} and the interior B° denote here $\partial B := \overline{B} \setminus B^\circ$ the edge of B .

Corollary 9.7 (Convergence of distribution functions). *Let $\mathbf{P}, \mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R})$ with distribution functions F, F_1, F_2, \dots . Then $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$ exactly if $F_n(x) \xrightarrow{n \rightarrow \infty} F(x)$ for all continuity points x of F .*

Proof. ' \Rightarrow ': If x is a continuity point of F , then $\mathbf{P}(\partial(-\infty; x]) = \mathbf{P}(\{x\}) = 0$. This means that – according to Theorem 9.6 (direction (i) \Rightarrow (v)) – that

$$F_n(x) = \mathbf{P}_n((-\infty; x]) \xrightarrow{n \rightarrow \infty} \mathbf{P}((-\infty; x]) = F(x).$$

' \Leftarrow ': According to Theorem 9.6 (direction (ii) \Rightarrow (i)), it suffices to show that $\mathbf{P}_n[f] \xrightarrow{n \rightarrow \infty} \mathbf{P}[f]$ for all bounded, Lipschitz functions f . Wlog, we assume that $|f| \leq 1$ and f has Lipschitz constant 1. For $\varepsilon > 0$ choose $N \in \mathbb{N}$ and continuity points $y_0 < \dots < y_N$ of F , so that $F(y_0) < \varepsilon$, $F(y_N) > 1 - \varepsilon$ and $y_i - y_{i-1} < \varepsilon$ for $i = 1, \dots, N$. Then $F_n(y_i) \xrightarrow{n \rightarrow \infty} F(y_i)$ and

$$f \leq 1_{(-\infty, y_0]} + 1_{(y_N, \infty)} + \sum_{i=1}^{N-1} (f(y_i) + \varepsilon) 1_{(y_i, y_{i+1}]},$$

as well as

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathbf{P}_n[f] &\leq \limsup_{n \rightarrow \infty} F_n(y_0) + 1 - F_n(y_N) + \sum_{i=1}^N (f(y_i) + \varepsilon)(F_n(y_i) - F_n(y_{i-1})) \\ &\leq 3\varepsilon + \sum_{i=1}^N f(y_i)(F(y_i) - F(y_{i-1})) \leq 4\varepsilon + \mathbf{P}[f]. \end{aligned}$$

With $\varepsilon \rightarrow 0$ and by replacing f with $1 - f$, we find $\mathbf{P}_n[f] \xrightarrow{n \rightarrow \infty} \mathbf{P}[f]$. \square

Remark 9.8 (The Theorem of deMoivre-Laplace). *In Example 9.3 we claimed that deMoivre-Laplace's Theorem makes a statement about weak convergence. The Theorem states that for $B(n, p)$ -distributed random variables X_n , $n = 1, 2, \dots$,*

$$\mathbf{P}\left(\frac{X_n - np}{\sqrt{np(1-p)}} \leq x\right) \xrightarrow{n \rightarrow \infty} \Phi(x),$$

where Φ is the distribution function of the standard normal distribution. As Corollary 9.7 shows, this means exactly the convergence in distribution to a standard normal distribution.

Corollary 9.9 (Slutzky's Theorem). *Let $X, X_1, X_2, \dots, Y_1, Y_2, \dots$ be random variables with values in E . If $X_n \xrightarrow{n \rightarrow \infty} X$ and $r(X_n, Y_n) \xrightarrow{n \rightarrow \infty}_p 0$, then $Y_n \xrightarrow{n \rightarrow \infty} X$.*

Proof. Let $f : E \rightarrow \mathbb{R}$ be bounded and Lipschitz-continuous with Lipschitz constant L . Then,

$$|f(x) - f(y)| \leq L \cdot r(x, y) \wedge (2\|f\|_\infty)$$

for all $x, y \in E$. From this,

$$\limsup_{n \rightarrow \infty} \mathbf{E}[f(X_n) - f(Y_n)] \leq \limsup_{n \rightarrow \infty} \mathbf{E}[L \cdot r(X_n, Y_n) \wedge (2\|f\|_\infty)] = 0$$

according to Lemma 7.5. Thus,

$$\limsup_{n \rightarrow \infty} |\mathbf{E}[f(Y_n)] - \mathbf{E}[f(X)]| \leq \limsup_{n \rightarrow \infty} |\mathbf{E}[f(Y_n)] - \mathbf{E}[f(X_n)]| + |\mathbf{E}[f(X_n)] - \mathbf{E}[f(X)]| = 0,$$

and the claimed convergence follows with Theorem 9.6. \square

Theorem 9.10 (Continuous mapping theorem). *Let E be separable, (E', r') another metric space and $\varphi : E \rightarrow E'$ measurable and $U_\varphi \subseteq E$ the set of discontinuity points of φ .*

1. *If $\mathbf{P}, \mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(E)$ and $\mathbf{P}(U_\varphi) = 0$ and $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$, then $\varphi_* \mathbf{P}_n \xrightarrow{n \rightarrow \infty} \varphi_* \mathbf{P}$.*
2. *If X, X_1, X_2, \dots are random variables with values in E and $\mathbf{P}(X \in U_\varphi) = 0$ and $X_n \xrightarrow{n \rightarrow \infty} X$, then also $\varphi(X_n) \xrightarrow{n \rightarrow \infty} \varphi(X)$.*

Proof. First, we note that 2. is an application of 1. if one sets $\mathbf{P}_n = (X_n)_* \mathbf{P}$. The set U_φ is Borel-measurable, since

$$U_\varphi^{\delta, \varepsilon} = \{x \in E : \exists y, z \in B_\delta(x), r'(\varphi(y), \varphi(z)) > \varepsilon\}$$

is Borel-measurable (here the separability of E is included) and

$$U_\varphi = \bigcup_{n=1}^{\infty} \bigcap_{k=1}^{\infty} U_\varphi^{1/k, 1/n}.$$

Let $G \subseteq E'$ be open and $x \in \varphi^{-1}(G) \cap U_\varphi^c$. Since φ is continuous in x , there is a $\delta > 0$ with $\varphi(y) \in G$ (i.e. $y \in \varphi^{-1}(G)$) for all y with $r(x, y) < \delta$. Therefore, $\varphi^{-1}(G) \cap U_\varphi^c \subseteq (\varphi^{-1}(G))^\circ$. This follows with Theorem 9.6 (direction (i) \Rightarrow (iii))

$$\begin{aligned} \varphi_* \mathbf{P}(G) &= \mathbf{P}(\varphi^{-1}(G)) = \mathbf{P}(\varphi^{-1}(G) \cap U_\varphi^c) \leq \mathbf{P}((\varphi^{-1}(G))^\circ) \\ &\leq \liminf_{n \rightarrow \infty} \mathbf{P}_n((\varphi^{-1}(G))^\circ) \leq \liminf_{n \rightarrow \infty} \mathbf{P}_n(\varphi^{-1}(G)) = \liminf_{n \rightarrow \infty} \varphi_* \mathbf{P}_n(G). \end{aligned}$$

Again due to theorem 9.6 (direction (iii) \Rightarrow (i)), this implies $\varphi_* \mathbf{P}_n \xrightarrow{n \rightarrow \infty} \varphi_* \mathbf{P}$. □

Apart from the vague convergence, convergence in distribution is the weakest form of convergence. However, there is a connection with almost sure convergence, as the following theorem shows.

Theorem 9.11 (Weak and almost sure convergence, Skorohod). *Let X, X_1, X_2, \dots be random variables with values in a complete and separable space (E, r) . Then, $X_n \xrightarrow{n \rightarrow \infty} X$ holds if and only if there is a probability space on which random variables Y, Y_1, Y_2, \dots are defined with $Y_n \xrightarrow{n \rightarrow \infty} Y$ *f.s.* and $Y \stackrel{d}{=} X, Y_1 \stackrel{d}{=} X_1, Y_2 \stackrel{d}{=} X_2, \dots$*

Proof. ' \Leftarrow ': This is clear, since almost sure convergence implies weak convergence (see Proposition 9.5).

' \Rightarrow ': We extend the probability space on which X is defined, and we set $Y = X$. Let $E = \{1, \dots, m\}$ be finite, U be uniformly distributed on $[0, 1]$ and independent of Y , and W_1, W_2, \dots independent with

$$\mathbf{P}(W_n = k) = \frac{\mathbf{P}(X_n = k) - \mathbf{P}(X = k) \wedge \mathbf{P}(X_n = k)}{1 - \sum_{l=1}^m \mathbf{P}(X = l) \wedge \mathbf{P}(X_n = l)}.$$

We set $Y_n = k$ if either

$$X = k \text{ and } U \leq \frac{\mathbf{P}(X_n = k)}{\mathbf{P}(X = k)}$$

or

$$X = l \text{ and } U > \frac{\mathbf{P}(X_n = l)}{\mathbf{P}(X = l)} \text{ and } W_n = k.$$

Then

$$\begin{aligned} \mathbf{P}(Y_n = k) &= \mathbf{P}(X = k) \cdot \frac{\mathbf{P}(X_n = k)}{\mathbf{P}(X = k)} \wedge 1 \\ &+ \sum_{l=1}^m \mathbf{P}(X = l) \cdot \left(1 - \frac{\mathbf{P}(X_n = l)}{\mathbf{P}(X = l)}\right) \frac{\mathbf{P}(X_n = k) - \mathbf{P}(X = k) \wedge \mathbf{P}(X_n = k)}{1 - \sum_{l'=1}^m \mathbf{P}(X = l') \wedge \mathbf{P}(X_n = l')} \\ &= \mathbf{P}(X_n = k) \wedge \mathbf{P}(X = k) \\ &+ \sum_{l=1}^m (\mathbf{P}(X = l) - \mathbf{P}(X_n = l) \wedge \mathbf{P}(X = l)) \frac{\mathbf{P}(X_n = k) - \mathbf{P}(X = k) \wedge \mathbf{P}(X_n = k)}{1 - \sum_{l'=1}^m \mathbf{P}(X = l') \wedge \mathbf{P}(X_n = l')} \\ &= \mathbf{P}(X_n = k). \end{aligned}$$

Thus $Y_n \stackrel{d}{=} X_n$. Since according to the condition $\mathbf{P}(X_n = k) \xrightarrow{n \rightarrow \infty} \mathbf{P}(X = k)$, the almost sure convergence follows.

For general E , let $p = 1, 2, \dots$ and choose a partition of E in sets B_1, B_2, \dots in E with $\mathbf{P}(Y \in \partial B_k) = 0$ and diameter at most 2^{-p} . Choose m large enough, so that $\mathbf{P}(Y \notin B_0) < 2^{-p}$ with $B_0 := E \setminus \bigcup_{k \leq m} B_k$. For $k = 1, 2, \dots$, define random variables $\tilde{Z}, \tilde{Z}_1, \tilde{Z}_2, \dots$ such that $\tilde{Z} = k$ exactly when $Y \in B_k$ and $\tilde{Z}_n = k$ if $Y_n \in B_k$. Then $\tilde{Z}_n \xrightarrow{n \rightarrow \infty} \tilde{Z}$. Since $\tilde{Z}, \tilde{Z}_1, \dots$ only takes values in a finite set, we can use random variables Z, Z_1, Z_2, \dots with $Z_n \xrightarrow{n \rightarrow \infty}_{f.s.} Z$. Furthermore, let $W_{n,k}$ be random variables with distribution $\mathbf{P}[X_n \in \cdot | X_n \in B_k]$ and $\tilde{Y}_{n,p} = \sum_k W_{n,k} 1_{Z_n=k}$, so that $\tilde{Y}_{n,p} \stackrel{d}{=} X_n$ for all n . It is now clear

$$\left\{r(\tilde{Y}_{n,p}, Y) > 2^{-p}\right\} \subseteq \{Z_n \neq Z\} \cup \{Y \in B_0\}.$$

Since $Z_n \xrightarrow{n \rightarrow \infty}_{f.s.} Z$ and $\mathbf{P}\{Y \in B_0\} < 2^{-p}$, for each p there are numbers $n_1 < n_2 < \dots$ with

$$\mathbf{P}\left(\bigcup_{n \geq n_p} \{r(\tilde{Y}_{n,p}, Y) > 2^{-p}\}\right) < 2^{-p}$$

for all p . With the Borel-Cantelli lemma we get

$$\sup_{n \geq n_p} r(\tilde{Y}_{n,p}, Y) \leq 2^{-p}$$

for almost all p . We therefore define $Y_n := \tilde{Y}_{n,p}$ for $n_p \leq n < n_{p+1}$ and note that $X_n \stackrel{d}{=} Y_n \xrightarrow{n \rightarrow \infty}_{f.s.} Y$. \square

9.2 Prohorov' Theorem

In this section, we first examine the concept of vague convergence. We will restrict ourselves to the space $E = \mathbb{R}$. (Most of the statements shown here are still valid in locally compact spaces). It is already clear that weak convergence of distributions implies vague convergence (since all continuous functions with compact support are bounded), and that the weak convergence is

equivalent to the convergence of the distribution functions (Corollary 9.7). The main result here is the theorem of Helly (Theorem 9.13), which states that every sequence of probability measures has a vaguely convergent subsequence.

We then examine the question when a sequence of probability measures also has weakly convergent subsequence. This leads us to the notion of tightness of probability measures and Prohorov's theorem (Theorem 9.19).

As we have already seen in Remark 9.2.1, it can be that the vague limit measure of probability measures is not a probability measure. However, the following result shows that the limit measure has total mass at most 1.

Lemma 9.12 (Mass loss at vague convergence). *Let $\mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R})$ and μ a measure on $\mathcal{B}(\mathbb{R})$ with $\mathbf{P}_n[f] \xrightarrow{n \rightarrow \infty} \mu[f]$, $f \in \mathcal{C}_c(\mathbb{R})$, then $\mu \in \mathcal{P}_{\leq 1}(\mathbb{R})$ applies.*

Proof. Let $f_1, f_2, \dots \in \mathcal{C}_c(\mathbb{R})$ with $f_k \uparrow 1$. Then with monotonic convergence

$$\mu(\mathbb{R}) = \sup_{k \in \mathbb{N}} \mu[f_k] = \sup_{k \in \mathbb{N}} \limsup_{n \rightarrow \infty} \mathbf{P}_n[f_k] \leq 1.$$

□

Theorem 9.13 (Helly's theorem). *Let $\mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R})$. Then there is a subsequence $(n_k)_{k=1,2,\dots}$ and a $\mu \in \mathcal{P}_{\leq 1}(\mathbb{R})$ with $\mathbf{P}_{n_k} \xrightarrow[k \rightarrow \infty]{v} \mu$.*

Proof. Let F_1, F_2, \dots be the distribution functions of $\mathbf{P}_1, \mathbf{P}_2, \dots$. Further, let (x_1, x_2, \dots) be a count of \mathbb{Q} . Since $[0, 1]$ is compact, for each sequence there is $(F_n(x_i))_{n=1,2,\dots}$ a convergent subsequence. By means of a diagonal argument, there is a sequence $(n_k)_{k=1,2,\dots}$ such that $(F_{n_k}(x_i))_{k=1,2,\dots}$ for all i against a limit $G(x_i)$ converges to \mathbb{Q} . We define

$$F(x) := \inf\{G(r) : r \in \mathbb{Q}, r > x\}.$$

Since all F_n and therefore G have non-negative increments, the same applies to F . From the definition of F and the monotonicity of G , it also follows that F is right-continuous. According to Proposition 2.19, there is a measure μ on \mathbb{R} with $\mu((x, y]) = F(y) - F(x)$ for all $x, y \in \mathbb{R}, x \leq y$. It remains to show that $\mathbf{P}_n[f] \xrightarrow{n \rightarrow \infty} \mu[f]$ for all $f \in \mathcal{C}_c(\mathbb{R})$. Wlog we can assume that $f \geq 0$.

It is $F_n(x) \xrightarrow{n \rightarrow \infty} F(x)$ at all continuity points x of F by construction. There is a countable set $D \subseteq \mathbb{R}$ such that F is continuous on D^c is continuous. This means that $\mathbf{P}_n(U) \xrightarrow{n \rightarrow \infty} \mu(U)$ for all finite unions U of intervals with vertices in D^c . Now let $B \subseteq \mathbb{R}$ be open and bounded. Let U_1, U_2, \dots and V_1, V_2, \dots be sequences of finite unions of open intervals with vertices in D^c such that $U_k \uparrow B, V_k \downarrow \bar{B}$. Then,

$$\begin{aligned} \mu(B) &= \lim_{k \rightarrow \infty} \mu(U_k) = \lim_{k \rightarrow \infty} \liminf_{n \rightarrow \infty} \mathbf{P}_n(U_k) \leq \liminf_{n \rightarrow \infty} \mathbf{P}_n(B) \\ &\leq \limsup_{n \rightarrow \infty} \mathbf{P}_n(B) \leq \lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbf{P}_n(V_k) = \lim_{k \rightarrow \infty} \mu(V_k) = \mu(\bar{B}). \end{aligned}$$

Since $\mu(f = t) > 0$ for at most countably many t , and since $\mathbf{P}_n(f > t) \leq 1_{t \geq \|f\|}$, it follows with dominated convergence

$$\begin{aligned} \mu[f] &= \int_0^\infty \mu(f > t) dt \leq \int_0^\infty \liminf_{n \rightarrow \infty} \mathbf{P}_n(f > t) dt = \liminf_{n \rightarrow \infty} \int_0^\infty \mathbf{P}_n(f > t) dt = \liminf_{n \rightarrow \infty} \mathbf{E}_n[f] \\ &\leq \limsup_{n \rightarrow \infty} \mathbf{P}_n[f] = \limsup_{n \rightarrow \infty} \int_0^\infty \mathbf{P}_n(f > t) dt = \int_0^\infty \limsup_{n \rightarrow \infty} \mathbf{P}_n(f > t) dt \leq \int_0^\infty \mu(f \geq t) \\ &= \mu[f]. \end{aligned}$$

□

We now return to the case of a general metric space (E, r) . To show the existence of accumulation points in the sense of weak convergence, it must be ensured that limit measures are again limit measures are again probability measures. In particular no mass is lost at the boundary crossing as in the case of vague convergence (see Lemma 9.12). Here, the concept of *tightness* is central.

Definition 9.14 (Tightness). *Let \mathcal{K} be the system of all compact sets in E . A family $(\mathbf{P}_i)_{i \in I}$ in $\mathcal{P}(E)$ is tight, if*

$$\sup_{K \in \mathcal{K}} \inf_{i \in I} \mathbf{P}_i(K) = 1.$$

A family $(X_i)_{i \in I}$ of E -valued random variables is tight if $((X_i)_ \mathbf{P})_{i \in I}$ is tight, i.e.*

$$\sup_{K \in \mathcal{K}} \inf_{i \in I} \mathbf{P}(X_i \in K) = 1.$$

Remark 9.15 (Equivalent formulations). *1. The definition of the tightness of a family $(\mathbf{P}_i)_{i \in I}$ in $\mathcal{P}(E)$ is equivalent to the following condition: for all $\varepsilon > 0$ there exists $K \subseteq E$ compact with $\inf_{i \in I} \mathbf{P}_i(K) \geq 1 - \varepsilon$.*

2. If $E = \mathbb{R}^d$, a family $(\mathbf{P}_i)_{i \in I}$ is tight if and only if

$$\sup_{r > 0} \inf_{i \in I} \mathbf{P}_i(B_r(0)) = 1,$$

where $B_r(0)$ is the sphere around 0 with radius r .

3. In Lemma 2.9 we have shown that $\mathbf{P} \in \mathcal{P}(E)$ is tight if (E, r) is complete and is separable. It also follows that every finite family of probability measures on the Borel's σ -algebra of a Polish space is tight.

4. Further, a countable family $(\mathbf{P}_i)_{i=1,2,\dots}$ is of probability measures on a Polish space (E, r) is tight if and only if

$$\sup_{K \in \mathcal{K}} \liminf_{i=1,2,\dots} \mathbf{P}_i(K) = 1.$$

Proof. ' \Rightarrow ': This is clear, since $\liminf_{i=1,2,\dots} \mathbf{P}_i(K) \geq \inf_{i=1,2,\dots} \mathbf{P}_i(K) = 1$.

' \Leftarrow ': Let $\varepsilon > 0$ and K such that $\liminf_{i=1,2,\dots} \mathbf{P}_i(K) \geq 1 - \varepsilon/2$. Choose N such that $\inf_{i=N+1, N+2, \dots} \mathbf{P}_i(K) \geq 1 - \varepsilon$ and K_1, \dots, K_N compact such that $\mathbf{P}_i(K_i) \geq 1 - \varepsilon$ for $i = 1, \dots, N$. Since $\tilde{K} = K \cup K_1 \cup \dots \cup K_N$ is compact and $\inf_{i=1,2,\dots} \mathbf{P}_i(\tilde{K}) \geq 1 - \varepsilon$ the tightness of $(\mathbf{P}_i)_{i=1,2,\dots}$ follows. □

Example 9.16 (Tight sets of probability measures). *1. If E is compact, every family of probability measures on $\mathcal{B}(E)$ is tight.*

2. A family $(X_i)_{i \in I}$ of real-valued random variables with

$$\sup_{i \in I} \mathbf{P}[|X_i|] < \infty,$$

is tight. This is because

$$\inf_{r > 0} \sup_{i \in I} \mathbf{P}(|X_i| \geq r) \leq \inf_{r > 0} \sup_{i \in I} \frac{\mathbf{P}[|X_i|]}{r} = 0.$$

3. The family $(\delta_n)_{n=1,2,\dots}$, where δ_n is the Dirac measure on n , is not tight.

Lemma 9.17 (Vague convergence and tightness). *Let $\mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R})$ and $\mu \in \mathcal{P}_{\leq 1}(\mathbb{R})$ with*

$$\mathbf{P}_n \xrightarrow[v]{n \rightarrow \infty} \mu.$$

Then

$$\mu(\mathbb{R}) = 1 \quad \iff \quad (\mathbf{P}_n)_{n=1,2,\dots} \text{ is tight}$$

. In this case, $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mu$.

Proof. For $r > 0$ choose a $g_r \in \mathcal{C}_c(\mathbb{R})$, $1_{B_r(0)} \leq g_r \leq 1_{B_{r+1}(0)}$. Then $(\mathbf{P}_n)_{n=1,2,\dots}$ is tight if and only if

$$\sup_{r>0} \liminf_{n \rightarrow \infty} \mathbf{P}_n[g_r] = 1.$$

' \Rightarrow ': Since μ is continuous from below, we find

$$1 = \sup_{r>0} \mu(B_r(0)) \leq \sup_{r>0} \mu[g_r] = \sup_{r>0} \liminf_{n \rightarrow \infty} \mathbf{P}_n[g_r] \leq 1.$$

' \Leftarrow ': Let $(\mathbf{P}_n)_{n=1,2,\dots}$ be tight. Then, from Lemma 9.12,

$$1 \geq \mu(\mathbb{R}) = \sup_{r>0} \mu(B_r(0)) = \sup_{r>0} \mu[g_r] = \sup_{r>0} \liminf_{n \rightarrow \infty} \mathbf{P}_n[g_r] = 1.$$

It remains to show the weak convergence. Assuming that $(\mathbf{P}_n)_{n=1,2,\dots}$ is tight and $f \in \mathcal{C}_b(\mathbb{R})$. Then,

$$\begin{aligned} \limsup_{n \rightarrow \infty} |\mathbf{P}_n[f] - \mu[f]| &\leq \inf_{r>0} \limsup_{n \rightarrow \infty} (|\mathbf{P}_n[f - fg_r]| + |\mathbf{P}_n[fg_r] - \mu[fg_r]| + |\mu[f - fg_r]|) \\ &\leq \|f\| \inf_{r>0} \limsup_{n \rightarrow \infty} \mathbf{P}_n(B_r(0)^c) + \inf_{r>0} \mu[B_r(0)^c] = 0, \end{aligned}$$

and $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mu$ follows. □

Corollary 9.18 (Weak convergence and tightness). *Let $\mathbf{P}, \mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R})$. If $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$, then $(\mathbf{P}_n)_{n \in \mathbb{N}}$ is tight.*

Proof. Since weak convergence of $\mathbf{P}_1, \mathbf{P}_2, \dots$ to \mathbf{P} implies vague convergence, for $\mathbf{P}, \mathbf{P}_1, \mathbf{P}_2, \dots$ the conditions of Lemma 9.17 and $\mathbf{P}(\mathbb{R}) = 1$ are satisfied. Therefore, $(\mathbf{P}_n)_{n \in \mathbb{N}}$ is tight. □

To determine the weak convergence of probability measures Theorem 9.6 is helpful. We now turn to the question whether a sequence of probability measures can have an accumulation point. This means that there is a subsequence that converges weakly to a probability measure.

Theorem 9.19 (Prohorov's theorem). *Let (E, r) be complete and separable and $(\mathbf{P}_i)_{i \in I}$ a family in $\mathcal{P}(E)$. The following are equivalent:*

1. The family $(\mathbf{P}_i)_{i \in I}$ is relatively compact with respect to the topology of weak convergence, i.e. every sequence in $(\mathbf{P}_i)_{i \in I}$ has a weakly convergent subsequence.

2. For every $\varepsilon > 0$ there is an $N \in \mathbb{N}$ and $x_1, \dots, x_N \in E$, so that

$$\inf_{i \in I} \mathbf{P}_i \left(\bigcup_{k=1}^N B_\varepsilon(x_k) \right) \geq 1 - \varepsilon.$$

3. The family $(\mathbf{P}_i)_{i \in I}$ is tight.

Proof. Let x_1, x_2, \dots be a dense subsequence in E (which exists since (E, r) is separable).

1. \Rightarrow 2.: Suppose 2. is not true. Then there is $\varepsilon > 0$ and for each $N = 1, 2, \dots$ a \mathbf{P}_{i_N} with $\mathbf{P}_{i_N} \left(\bigcup_{k=1}^N B_\varepsilon(x_k) \right) \leq 1 - \varepsilon$. By relative compactness, there would then be some subsequence $(\mathbf{P}_{i_M})_{M=1,2,\dots}$ which is weakly convergent to some $\mathbf{P} \in \mathcal{P}(E)$. Thus, because of Theorem 9.6 ((i) \Rightarrow (iii)) we find that

$$1 = \mathbf{P}(E) = \sup_{N \in \mathbb{N}} \mathbf{P} \left(\bigcup_{k=1}^N B_\varepsilon(x_k) \right) \leq \sup_{N \in \mathbb{N}} \liminf_{M \rightarrow \infty} \mathbf{P}_{i_M} \left(\bigcup_{k=1}^N B_\varepsilon(x_k) \right) \leq 1 - \varepsilon,$$

thus a contradiction.

2. \Rightarrow 3.: Let $\varepsilon > 0$. For $j = 1, 2, \dots$ we choose x_{j1}, \dots, x_{jN_j} such that

$$\inf_{i \in I} \mathbf{P}_i \left(\bigcup_{k=1}^{N_j} B_{\varepsilon 2^{-j}}(x_{jk}) \right) > 1 - \varepsilon 2^{-j}.$$

We further set

$$K := \bigcap_{j=1}^{\infty} \bigcup_{k=1}^{N_j} B_{\varepsilon 2^{-j}}(x_{jk}).$$

Then $K \subseteq E$ is totally bounded by construction, according to Proposition A.9 therefore relatively compact, so \overline{K} is compact. Furthermore

$$\sup_{i \in I} \mathbf{P}_i(\overline{K}^c) \leq \sup_{i \in I} \sum_{j=1}^{\infty} \mathbf{P}_i \left(\bigcap_{k=1}^{N_j} (B_{\varepsilon 2^{-j}}(x_{jk}))^c \right) \leq \varepsilon.$$

Thus the family $(\mathbf{P}_i)_{i \in I}$ is tight.

3. \Rightarrow 1.: Let $\mathbf{P}_1, \mathbf{P}_2, \dots$ be a sequence in the family of the family $(\mathbf{P}_i)_{i \in I}$. The aim is to determine a convergent subsequence. For this purpose, we choose compact sets $K_1 \subseteq K_2 \subseteq \dots \subseteq E$ with $\inf_{n=1,2,\dots} \mathbf{P}_n(K_j) \geq 1 - 1/j$. Further, we choose the system of compact sets

$$\mathcal{K} := \left\{ \bigcup_{k=1}^N K_{j_k} \cap \overline{B_{\varepsilon_k}(x_k)} : N, j_k \in \mathbb{N}, \varepsilon_k \in \mathbb{Q}^+ \right\}.$$

Thus \mathcal{K} is a semi-ring and also a \cap -stable generator of $\sigma(E)$. Since \mathcal{K} is countable, we can create a subsequence $\mathbf{P}_{n_1}, \mathbf{P}_{n_2}, \dots$ from $\mathbf{P}_1, \mathbf{P}_2, \dots$ so that $\mathbf{P}_{n_k}(A)$ converges for all $A \in \mathcal{K}$. We set for $A \in \mathcal{K}$

$$\mu(A) = \lim_{k \rightarrow \infty} \mathbf{P}_{n_k}(A).$$

Thus, according to theorem 2.10 and Theorem ??, the content μ on \mathcal{K} uniquely becomes a measure on $\sigma(\mathcal{K}) = \mathcal{B}(E)$ can be continued. The thus defined μ is a probability measure because

$$1 \geq \mu(E) = \sup_{j=1,2,\dots} \mu(K_j) = \sup_{j=1,2,\dots} \lim_{k \rightarrow \infty} \mathbf{P}_{n_k}(K_j) \geq \sup_{j=1,2,\dots} 1 - 1/j = 1.$$

Furthermore, for every open set A

$$\mu(A) = \sup_{A \supseteq H \in \mathcal{K}} \mu(H) = \sup_{A \supseteq H \in \mathcal{K}} \lim_{k \rightarrow \infty} \mathbf{P}_{n_k}(H) \leq \liminf_{k \rightarrow \infty} \mathbf{P}_{n_k}(A).$$

Now follows $\mathbf{P}_{n_k} \xrightarrow{k \rightarrow \infty} \mu$ from theorem 9.6 ((iii) \Rightarrow (i)). □

9.3 Separating classes of functions

Now we will introduce separating classes of functions. In particular, this will shed some light on the usefulness of characteristic functions and Laplace transforms of distributions (see Definition 6.11). These are based on two specific classes of functions that are separating.

Definition 9.20 (Classes of functions separating points and separating function classes).

1. A function class $\mathcal{M} \subseteq \mathcal{C}(E)$ is said to separate points in E if for all $x, y \in E$ with $x \neq y$ there exists an $f \in \mathcal{M}$ with $f(x) \neq f(y)$.
2. A class of functions $\mathcal{M} \subseteq \mathcal{C}(E)$ is called separating in $\mathcal{P}(E)$ if from $\mathbf{P}, \mathbf{Q} \in \mathcal{P}(E)$ and

$$\mathbf{P}[f] = \mathbf{Q}[f] \text{ for all } f \in \mathcal{M}$$

implies that $\mathbf{P} = \mathbf{Q}$.

Example 9.21. 1. The class of functions $\mathcal{M} := \mathcal{C}_b(E)$ is both, separating points and separating. Namely, if $x \neq y$, then $z \mapsto r(x, z) \wedge 1$ is a bounded, continuous function that separates x and y . Furthermore, if $\mathbf{P}, \mathbf{Q} \in \mathcal{P}(E)$ and $\mathbf{P} \neq \mathbf{Q}$, then there is an open ball A with $\mathbf{P}(A) \neq \mathbf{Q}(A)$. Let f_1, f_2, \dots be a sequence in $\mathcal{C}_b(E)$ with $f_n \uparrow 1_A$. If $\mathbf{P}[f_n] = \mathbf{Q}[f_n]$ for all $n = 1, 2, \dots$, then it would also

$$\mathbf{P}(A) = \lim_{n \rightarrow \infty} \mathbf{P}[f_n] = \lim_{n \rightarrow \infty} \mathbf{Q}[f_n] = \mathbf{Q}(A)$$

in contradiction to the assumption.

2. The class of functions $\{x \mapsto cx : c \in \mathbb{R}\}$ of all linear functions separates points, but is not separating.

The next result requires the Stone-Weierstrass theorem, which we repeat first.

Definition 9.22 (Algebra). A set system $\mathcal{M} \subseteq \mathcal{C}(E)$ is called an algebra, if $1 \in \mathcal{M}$, and if $\alpha, \beta \in \mathbb{R}$ and it contains f, g it also contains $\alpha f + \beta g$, as well as fg .

Theorem 9.23 (Stone-Weierstrass). Let (E, r) be compact and $\mathcal{M} \subseteq \mathcal{C}_b(E)$ an algebra separating points. Then, \mathcal{M} is dense in $\mathcal{C}_b(E)$ with respect to the supremum norm.

Proof. See some lecture on Analysis. □

Theorem 9.24 (Algebras separating points and separating algebras).

Let (E, r) be complete and separable. If $\mathcal{M} \subseteq \mathcal{C}_b(E)$ separates points and is such that $f, g \in \mathcal{M}$ implies $fg \in \mathcal{M}$. Then \mathcal{M} is separating.

Proof. Let $\mathbf{P}, \mathbf{Q} \in \mathcal{P}(E)$. Without restriction, $1 \in \mathcal{M}$, since $\mathbf{P}[1] = \mathbf{Q}[1]$ always holds. Thus \mathcal{M} is wlog an algebra. Let $\varepsilon > 0$ and K be compact such that $\mathbf{P}(K) > 1 - \varepsilon$, $\mathbf{Q}(K) > 1 - \varepsilon$. For $g \in \mathcal{C}_b(E)$, according to the Stone-Weierstrass Theorem 9.23 there is a sequence $(g_n)_{n=1,2,\dots}$ in \mathcal{M} with

$$\sup_{x \in K} |g_n(x) - g(x)| \xrightarrow{n \rightarrow \infty} 0. \quad (9.4)$$

Now,

$$\begin{aligned} |\mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{Q}[ge^{-\varepsilon g^2}]| &\leq |\mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{P}[ge^{-\varepsilon g^2}; K]| \\ &\quad + |\mathbf{P}[ge^{-\varepsilon g^2}; K] - \mathbf{P}[g_n e^{-\varepsilon g_n^2}; K]| \\ &\quad + |\mathbf{P}[g_n e^{-\varepsilon g_n^2}; K] - \mathbf{P}[g_n e^{-\varepsilon g_n^2}]| \\ &\quad + |\mathbf{P}[g_n e^{-\varepsilon g_n^2}] - \mathbf{Q}[g_n e^{-\varepsilon g_n^2}]| \\ &\quad + |\mathbf{Q}[g_n e^{-\varepsilon g_n^2}] - \mathbf{Q}[g_n e^{-\varepsilon g_n^2}; K]| \\ &\quad + |\mathbf{Q}[g_n e^{-\varepsilon g_n^2}] - \mathbf{Q}[ge^{-\varepsilon g^2}; K]| \\ &\quad + |\mathbf{Q}[ge^{-\varepsilon g^2}; K] - \mathbf{Q}[ge^{-\varepsilon g^2}]| \end{aligned}$$

We restrict the first term by

$$|\mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{P}[ge^{-\varepsilon g^2}; K]| \leq \frac{C}{\sqrt{\varepsilon}} \mathbf{P}(K^c) \leq C\sqrt{\varepsilon}$$

with $C = \sup_{x \geq 0} x e^{-x^2}$; analogous to the third, fifth and last terms. The second and penultimate terms converge to 0 for $n \rightarrow \infty$ due to (9.4). Since \mathcal{M} is an algebra, $g_n e^{-\varepsilon g_n^2}$ can be approximated by functions in \mathcal{M} , which means that the fourth term for $n \rightarrow \infty$ converges to 0. This means that

$$|\mathbf{P}[g] - \mathbf{Q}[g]| = \lim_{\varepsilon \rightarrow 0} |\mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{Q}[ge^{-\varepsilon g^2}]| \leq 4C \lim_{\varepsilon \rightarrow 0} \sqrt{\varepsilon} = 0.$$

Since g was arbitrary and $\mathcal{C}_b(E)$ is separating, the following follows $\mathbf{P} = \mathbf{Q}$. \square

We now come back to the characteristic function and the Laplace transform back. As already mentioned, the usefulness of the characteristic function and the Laplace transforms is due to the fact that they are distribution-determining.

Proposition 9.25 (Characteristic function distribution-determining).

A probability measure $\mathbf{P} \in \mathcal{P}(\mathbb{R}^d)$ ($\mathbf{P} \in \mathcal{P}(\mathbb{R}_+^d)$) is uniquely characterized by the characteristic function $\psi_{\mathbf{P}}$ (the Laplace transform $\mathcal{L}_{\mathbf{P}}$).

Proof. We show the statement only for characteristic functions that are proven for Laplace transforms is proven analogously. We establish that the set $\mathcal{M} := \{x \mapsto e^{itx}; t \in \mathbb{R}^d\}$ in \mathbb{R}^d separates points. Since $\mathcal{M} \subseteq \mathcal{C}_b(\mathbb{R}^d)$ and is closed under product formation, it is also separating according to theorem 9.24. This is shown by the statement. \square

Corollary 9.26 (Independence and characteristic function). *1. A family $(X_j)_{j \in I}$ of real-valued random variables is independent if and only if for all $J \subseteq_f I$*

$$\mathbf{E} \left[\prod_{j \in J} e^{it_j X_j} \right] = \prod_{j \in J} \mathbf{E} [e^{it_j X_j}] \quad (9.5)$$

for all $(t_j)_{j \in J} \in \mathbb{R}^J$ is valid.

2. A family $(X_j)_{j \in I}$ of random variables with values in \mathbb{R}_+ is independent if and only if for all $J \subseteq_f I$

$$\mathbf{E} \left[\prod_{j \in I} e^{-t_j X_j} \right] = \prod_{j \in J} \mathbf{E} [e^{-t_j X_j}]$$

for all $(t_j)_{j \in J} \in \mathbb{R}^J$ applies.

Proof. We only show the first statement, the second follows analogously. If $(X_j)_{j \in I}$ is independent, then according to Lemma 8.4 the random variables $(e^{it_j X_j})_{j \in I}$ for all $(t_j)_{j \in J} \in \mathbb{R}^J$ independently. Thus (9.5) follows from Proposition 9.25. Conversely, the following applies (9.5). On the one hand, the left-hand side of (9.5) represents the characteristic function of the distribution $((X_j)_{j \in J})_* \mathbf{P}$. On the other hand, the right side of (9.5) is the characteristic function of $\otimes_{j \in J} (X_j)_* \mathbf{P}$. Since the characteristic function according to Proposition 9.25 is the joint distribution of $(X_j)_{j \in J}$ is uniquely determined, $((X_j)_{j \in J})_* \mathbf{P} = \otimes_{j \in J} (X_j)_* \mathbf{P}$. The independence of $(X_j)_{j \in I}$ thus follows from Proposition 8.2. \square

9.4 The theorem of Lévy

We now want to analyze the relationship between weak convergence and the convergence of the characteristic functions of the underlying distributions. Let $\mathbf{P}, \mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R}^d)$. How to get from Proposition 9.27, the weak convergence follows $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$ follows from the pointwise convergence of the characteristic functions, $\psi_{\mathbf{P}_n}(t) \xrightarrow{n \rightarrow \infty} \psi_{\mathbf{P}}(t)$, $t \in \mathbb{R}^d$, given $(\mathbf{P}_n)_{n \in \mathbb{N}}$ is tight. The decisive factor is that the tightness of the family $(\mathbf{P}_n)_{n \in \mathbb{N}}$ can also be read from the characteristic functions as we will show in Proposition 9.32 will show. This leads to the statement of Lévy's continuity theorem (Theorem 9.33), which states when the pointwise limit of characteristic functions is again a characteristic function of a probability measure.

Proposition 9.27 (Separating class of functions and weak convergence). *Let (E, r) be complete and separable and $\mathbf{P}, \mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(E)$. Then are equivalent:*

1. $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$.

2. $(\mathbf{P}_n)_{n=1,2,\dots}$ is tight and there is a separating family $\mathcal{M} \subseteq C_b(E)$ with

$$\mathbf{P}_n[f] \xrightarrow{n \rightarrow \infty} \mathbf{P}[f] \text{ for all } f \in \mathcal{M}.$$

Proof. 1. \rightarrow 2. According to corollary 9.18 is $(\mathbf{P}_n)_{n=1,2,\dots}$ is tight. The second part of 2. applies because of the definition of weak convergence.

2. \Rightarrow 1. Suppose $(\mathbf{P}_n)_{n=1,2,\dots}$ is tight and $\mathbf{P}_1, \mathbf{P}_2, \dots$ does not converge weakly against \mathbf{P} . Then there is $\varepsilon > 0$, a $f \in \mathcal{C}_b(E)$ and a subsequence $(n_k)_{k=1,2,\dots}$ such that

$$(9.6)$$

$$|\mathbf{P}_{n_k}[f] - \mathbf{P}[f]| > \varepsilon \text{ for all } k. \quad (9.7)$$

According to theorem 9.19 there is a subsequence $(n_{k_\ell})_{\ell=1,2,\dots}$ and a $\mathbf{Q} \in \mathcal{P}(E)$, such that $\mathbf{P}_{n_{k_\ell}} \xrightarrow{\ell \rightarrow \infty} \mathbf{Q}$. Because of (9.6) is

$$|\mathbf{P}[f] - \mathbf{Q}[f]| \geq |\liminf_{\ell \rightarrow \infty} (\mathbf{P}[f] - \mathbf{P}_{n_{k_\ell}}[f]) + \liminf_{\ell \rightarrow \infty} (\mathbf{P}_{n_{k_\ell}}[f] - \mathbf{Q}[f])| > \varepsilon,$$

in particular therefore $\mathbf{P} \neq \mathbf{Q}$. On the other hand, for all $g \in \mathcal{M}$ we have

$$\mathbf{P}[g] = \lim_{\ell \rightarrow \infty} \mathbf{P}_{n_{k_\ell}}[g] = \mathbf{Q}[g].$$

Since \mathcal{M} is separating, this is a contradiction and 1. is shown. \square

Let $\mathbf{P} \in \mathcal{P}(\mathbb{R})$ and $\psi_{\mathbf{P}}$ be its characteristic function. We first show an estimation, which is important to relate tightness and the $\psi_{\mathbf{P}}$. bring.

Lemma 9.28 (tightness and the characteristic function). *Let $\mathbf{P} \in \mathcal{P}(\mathbb{R})$. Then for all $r > 0$*

$$\mathbf{P}((-\infty; -r] \cup [r; \infty)) \leq \frac{r}{2} \int_{-2/r}^{2/r} (1 - \psi_{\mathbf{P}}(t)) dt, \quad (9.8)$$

Proof. It is $\sin(x)/x \leq 1$ for $x \leq 2$ and $\sin x \leq x/2$ for $x \geq 2$. Let X be a random variable with distribution \mathbf{P} . Therefore, for every $c > 0$ according to Fubini

$$\begin{aligned} \int_{-c}^c (1 - \psi_{\mathbf{P}}(t)) dt &= \mathbf{P} \left[\int_{-c}^c (1 - e^{itX}) dt \right] = \mathbf{P} \left[2c - \frac{1}{iX} e^{itX} \Big|_{t=-c}^c \right] \\ &= 2c \mathbf{P} \left[1 - \frac{\sin(cX)}{cX} \right] \\ &\geq 2c \mathbf{P} \left[1 - \frac{\sin(cX)}{cX}; |cX| \geq 2 \right] \\ &\geq c \cdot \mathbf{P}(|cX| \geq 2) = c \mathbf{P}((-\infty; -\frac{2}{c}] \cup [\frac{2}{c}; \infty)), \end{aligned}$$

and the assertion follows with $c = 2/r$. \square

Definition 9.29 (Continuity of the same degree). *We repeat a definition from calculus. A set $\mathcal{M} \subseteq \mathcal{C}(\mathbb{R}^d)$ is called in $x \in \mathbb{R}^d$ is continuous to the same degree if*

$$\sup_{f \in \mathcal{M}} |f(y) - f(x)| \xrightarrow{y \rightarrow x} 0.$$

Remark 9.30 (Equivalent condition for sequences). *If $\mathcal{M} = \{f_1, f_2, \dots\}$, then the condition*

$$\limsup_{n \rightarrow \infty} |f_n(y) - f_n(x)| \xrightarrow{y \rightarrow x} 0$$

is equivalent.

Lemma 9.31 (Equal degree of continuity and convergence). *Let $f_1, f_2, \dots \in \mathcal{C}(\mathbb{R}^d)$, so that $f_n \xrightarrow{n \rightarrow \infty} f$ pointwise for a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ holds. Exactly then f is continuous in 0, if $(f_n)_{n=1,2,\dots}$ is uniformly continuous in 0.*

Proof. If $(f_n)_{n=1,2,\dots}$ is uniformly continuous in 0, then

$$|f(t) - f(0)| = \left| \lim_{n \rightarrow \infty} (f_n(t) - f_n(0)) \right| \leq \limsup_{n \rightarrow \infty} |f_n(t) - f_n(0)| \xrightarrow{t \rightarrow 0} 0.$$

Conversely, if f is continuous is continuous in 0, then

$$\limsup_{n \rightarrow \infty} |f_n(t) - f_n(0)| \leq \limsup_{n \rightarrow \infty} |f_n(t) - f(t)| + |f(t) - f(0)| + |f(0) - f_n(0)| = |f(t) - f(0)| \xrightarrow{t \rightarrow 0} 0.$$

□

Proposition 9.32 (tightness and uniformity continuity). *Let $(\mathbf{P}_i)_{i \in I}$ be a family in $\mathcal{P}(\mathbb{R}^d)$. If $(\psi_{\mathbf{P}_i})_{i \in I}$ is uniformly continuous in 0, then $(\mathbf{P}_i)_{i \in I}$ is tight.*

Proof. It suffices to show that $((\pi_k)_* \mathbf{P}_i)_{i \in I}$ for all projections π_1, \dots, π_d is tight. The following obviously applies $\psi_{(\pi_k)_* \mathbf{P}_i}(t) = \psi_{\mathbf{P}_i}(te_k)$, if e_k is the k -th unit vector. It is therefore sufficient to prove the assertion in the case $d = 1$. Since $\psi_{\mathbf{P}_i}(0) = 1$ for all $i \in I$, we conclude from the continuity of the same degree, that

$$\sup_{i \in I} |1 - \psi_{\mathbf{P}_i}(t)| \xrightarrow{t \rightarrow 0} 0,$$

thus, see remark 9.15,

$$\begin{aligned} \sup_{r > 0} \inf_{i \in I} \mathbf{P}_i([-r; r]) &\geq 1 - \inf_{r > 0} \sup_{i \in I} \frac{r}{2} \int_{-2/r}^{2/r} (1 - \psi_{\mathbf{P}_i}(t)) dt \\ &\geq 1 - \inf_{r > 0} \frac{r}{2} \int_{-2/r}^{2/r} \sup_{i \in I} |1 - \psi_{\mathbf{P}_i}(t)| dt \\ &\geq 1 - 2 \inf_{r > 0} \sup_{t \in [0; 2/r]} \sup_{i \in I} |1 - \psi_{\mathbf{P}_i}(t)| = 1. \end{aligned}$$

This shows the assertion. □

Theorem 9.33 (Lévy's continuity theorem). *Let $\mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R}^d)$ and $\psi : \mathbb{R}^d \rightarrow \mathbb{C}$, so that $\psi_{\mathbf{P}_n}(t) \xrightarrow{n \rightarrow \infty} \psi(t)$ for all $t \in \mathbb{R}^d$. If ψ is continuous in 0, then $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$ for a $\mathbf{P} \in \mathcal{P}(\mathbb{R}^d)$ with $\psi_{\mathbf{P}} = \psi$.*

Proof. Since $\psi_{\mathbf{P}_n}$ converges pointwise to the function continuous in 0 ψ , it follows from Lemma 9.31 that $(\psi_{\mathbf{P}_n})_{n=1,2,\dots}$ is uniformly continuous in 0 is continuous. With Proposition 9.32 it follows that $(\mathbf{P}_n)_{n=1,2,\dots}$ is tight. Let $(n_k)_{k=1,2,\dots}$ be a subsequence and $\mathbf{P} \in \mathcal{P}(\mathbb{R}^d)$ such that $\mathbf{P}_{n_k} \xrightarrow{k \rightarrow \infty} \mathbf{P}$. Since $x \mapsto e^{itx}$ is a continuous, bounded function, it follows that $\psi_{\mathbf{P}_{n_k}}(t) \xrightarrow{k \rightarrow \infty} \psi_{\mathbf{P}}(t)$ for all $t \in \mathbb{R}^d$. On the other hand, according to condition, $\psi_{\mathbf{P}_n}(t) \xrightarrow{n \rightarrow \infty} \psi(t)$, from which $\psi_{\mathbf{P}} = \psi$ follows. This identifies ψ as a characteristic function of \mathbf{P} and since this uniquely determines \mathbf{P} , the following applies thus $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$. □

Example 9.34 (Theorem of deMoivre-Laplace). Let $S_n \sim B(n, p)$. The theorem of deMoivre-Laplace states that

$$S_n^* := \frac{S_n - np}{\sqrt{np(1-p)}} \xrightarrow{n \rightarrow \infty} N(0, 1). \quad (9.9)$$

We now want to show this again with the help of characteristic functions, i.e. $\psi_{S_n^*} \xrightarrow{n \rightarrow \infty} \psi_{N(0,1)}$ point by point. To do this, we use Proposition 6.12.3 and write with $q := 1 - p$ and $C_1, C_2, \dots \in \mathbb{C}$ with $\limsup_{n \rightarrow \infty} |C_n| < \infty$

$$\begin{aligned} \psi_{S_n^*}(t) &= \exp\left(-it\sqrt{\frac{np}{q}}\right) \cdot \psi_{B(n,p)}\left(\frac{t}{\sqrt{npq}}\right) \\ &= \exp\left(-it\sqrt{\frac{np}{q}}\right) \left(q + p \exp\left(\frac{it}{\sqrt{npq}}\right)\right)^n \\ &= \left(q \exp\left(-it\sqrt{\frac{p}{nq}}\right) + p \exp\left(it\sqrt{\frac{q}{np}}\right)\right)^n \\ &= \left(1 - qit\sqrt{\frac{p}{nq}} - q\frac{t^2}{2}\frac{p}{nq} + pit\sqrt{\frac{q}{np}} - p\frac{t^2}{2}\frac{q}{np} + \frac{C_n}{n^{3/2}}\right)^n \\ &= \left(1 - \frac{t^2}{2}\frac{1}{n} + \frac{C_n}{n^{3/2}}\right)^n \xrightarrow{n \rightarrow \infty} e^{-\frac{t^2}{2}} = \psi_{N(0,1)}(t). \end{aligned}$$

From theorem 9.33 now follows (9.9).

Lévy's continuity theorem can also be formulated with formulated with Laplace transforms. We state the theorem without proof without proof:

Theorem 9.35 (Lévy's continuity theorem for Laplace transforms). Let $\mathbf{P}_1, \mathbf{P}_2, \dots \in \mathcal{P}(\mathbb{R}_+^d)$ and $\mathcal{L} : \mathbb{R}^d \rightarrow [0, 1]$, so that $\mathcal{L}_{\mathbf{P}_n}(t) \xrightarrow{n \rightarrow \infty} \mathcal{L}(t)$ for all $t \in \mathbb{R}^d$. If \mathcal{L} is continuous in 0, then $\mathbf{P}_n \xrightarrow{n \rightarrow \infty} \mathbf{P}$ for a $\mathbf{P} \in \mathcal{P}(\mathbb{R}^d)$ with $\mathcal{L}_{\mathbf{P}} = \mathcal{L}$.

Example 9.36 (Convergence of the geometric to the exponential distribution). Let $X_n \sim \mu_{\text{geo}(p_n)}$ be distributed and $n \cdot p_n \xrightarrow{n \rightarrow \infty} \lambda$. Then

$$\begin{aligned} \mathcal{L}_{X_n/n}(t) &= \mathbf{P}[e^{-tX_n/n}] = \sum_{k=1}^{\infty} (1-p_n)^{k-1} p_n e^{-tk/n} \\ &= p_n e^{-t/n} \frac{1}{1 - (1-p_n)e^{-t/n}} \\ &= \frac{\lambda}{n(1 - (1 - \lambda/n)(1 - t/n))} + o(1/n) \\ &\xrightarrow{n \rightarrow \infty} \frac{\lambda}{\lambda + t}. \end{aligned}$$

Therefore, $\frac{X_n}{n} \xrightarrow{n \rightarrow \infty} Y$, where $Y \sim \mu_{\text{exp}(\lambda)}$, since

$$\mathcal{L}_{\text{exp}(\lambda)}(t) = \int_0^{\infty} \lambda e^{-\lambda a} e^{-ta} da = \frac{\lambda}{\lambda + t}.$$

10 limit sets in distribution

We will now apply our knowledge of weak convergence and characteristic functions in special situations. In section 10.1 we are concerned with statements about when the sum of random variables converges against a Poisson distributed random variable converges. In section 10.2 we will apply the central Lindeberg-Feller's central limit theorem, which provides a characterization for the weak convergence against a normal normal distribution. Section 10.3 finally deals with extensions for the case of multidimensional random variables. random variables.

10.1 Poisson convergence

We already know the statement that $B(n, p_n)$ for $n \cdot p_n \xrightarrow{n \rightarrow \infty} \lambda$ for large n weakly against the $\text{Poi}(\lambda)$ -distribution converges, see Example 10.1. In this section we generalize this statement, see theorem 10.5.

Example 10.1 (Poisson approximation of the binomial distribution). *Let $p_1, p_2, \dots \in [0, 1]$ be such that $n \cdot p_n \xrightarrow{n \rightarrow \infty} \lambda$. Then we already know from the stochastics lecture that*

$$B(n, p_n)(\{k\}) \xrightarrow{n \rightarrow \infty} \text{Poi}(\lambda)(\{k\}).$$

In other words, this is a statement about weak convergence:

$$B(n, p_n) \xrightarrow{n \rightarrow \infty} \text{Poi}(\lambda). \tag{10.1}$$

Lévy's theorem provides another way to prove this result. to prove this result. We recall the characteristic functions of the binomial and Poisson distribution from Example 6.13. We write directly

$$\begin{aligned} \psi_{B(n, p_n)}(t) &= \left(1 - p_n(1 - e^{it})\right)^n \\ &= \left(1 - \frac{n \cdot p_n}{n}(1 - e^{it})\right)^n \\ &\xrightarrow{n \rightarrow \infty} \exp(-\lambda(1 - e^{it})) = \psi_{\text{Poi}(\lambda)}(t). \end{aligned}$$

In particular, the characteristic functions of the binomial distributions converge pointwise to a function that is continuous in 0, namely the characteristic function of the Poisson distribution. From Theorem 9.33 follows (10.1).

In the following, we will see that the weak convergence against a Poisson distribution is even more general. For this we will use use generating end functions.

Remark 10.2 (generating end function). *Consider a random variable X with values in \mathbb{Z}_+ and define the generating function*

$$z \mapsto \varphi_X(z) := \mathbf{P}[z^X] = \sum_{k=0}^{\infty} z^k \mathbf{P}[X = k].$$

We note that for $z \in [0, 1]$ this is consistent with the Laplace transform of X is closely related because (with $z = e^{-t}$)

$$\mathcal{L}_X(t) = \mathbf{P}[e^{-tX}] = \mathbf{P}[z^X] = \varphi_X(z).$$

In particular, two properties of Laplace transforms to generating functions.

1. Generating functions determining the distribution, see Proposition 9.25: *The distribution of X is uniquely determined by $z \mapsto \varphi_X(z)$ for $z \in [0, 1]$.*
2. Weak convergence equivalent to the convergence of the generating functions, see theorem 9.33: *Let X_1, X_2, \dots a sequence of random variables with values in \mathbb{Z}_+ such that $\varphi_{X_n}(z) \xrightarrow{n \rightarrow \infty} \varphi(z)$ for $z \in [0, 1]$ for a function φ that is continuous in 1 is continuous from below. Then $X_n \xrightarrow{n \rightarrow \infty} X$ is for a random variable X with generating function φ .*

Sometimes generating functions are practical tools. By their definition, they are power series with radius of convergence $r \geq 1$. It is known that inside the radius of convergence, the derivative and sum are interchanged. So if $r > 1$, for example, we write

$$\varphi'_X(1) = \sum_{k=0}^{\infty} k z^{k-1} \mathbf{P}(X = k) \Big|_{z=1} = \sum_{k=0}^{\infty} k \mathbf{P}(X = k) = \mathbf{P}[X].$$

Analogous calculations for higher derivatives are also possible.

Definition 10.3 (Asymptotic negligibility). *A triangular family of random variables $(X_{nj})_{n=1,2,\dots,n,j=1,\dots,m_n}$ with $m_1, m_2, \dots \in \mathbb{N}$ is asymptotically negligible if for $n = 1, 2, \dots$ the random variables X_{n1}, \dots, X_{nm_n} are independent and*

$$\sup_{j=1,\dots,m_n} \mathbf{P}(|X_{nj}| > \varepsilon) \xrightarrow{n \rightarrow \infty} 0 \quad (10.2)$$

for all $\varepsilon > 0$. If $X_{ij} \geq 0$ for all i, j , then $m_n = \infty$ is also permitted.

Remark 10.4 (Equivalent formulation). *1. For a triangular family of random variables $(X_{nj})_{n=1,2,\dots,n,j=1,\dots,m_n}$ holds (10.2) exactly when*

$$\sup_{j=1,\dots,m_n} \mathbf{E}[|X_{nj}| \wedge 1] \xrightarrow{n \rightarrow \infty} 0.$$

2. Let $(X_{nj})_{n=1,2,\dots,n,j=1,\dots,m_n}$ be a triangular of \mathbb{Z}_+ -valued random variables. Then (10.2) exactly when

$$\inf_{z \in [0,1]} \inf_{j=1,\dots,m_n} \varphi_{X_{nj}}(z) = \inf_{j=1,\dots,m_n} \varphi_{X_{nj}}(0) = \inf_{j=1,\dots,m_n} \mathbf{P}(|X_{nj}| = 0) \xrightarrow{n \rightarrow \infty} 1. \quad (10.3)$$

Theorem 10.5 (Poisson convergence). *Let $(X_{nj})_{n=1,2,\dots,n,j=1,\dots,m_n}$ be a family of asymptotically negligible random variables with values in \mathbb{Z}_+ and $X \sim \text{Poi}(\lambda)$. Then applies*

$$\sum_{j=1}^{m_n} X_{nj} \xrightarrow{n \rightarrow \infty} X$$

exactly when holds:

1. $\sum_{j=1}^{m_n} \mathbf{P}(X_{nj} > 1) \xrightarrow{n \rightarrow \infty} 0$
2. $\sum_{j=1}^{m_n} \mathbf{P}(X_{nj} = 1) \xrightarrow{n \rightarrow \infty} \lambda.$

We prepare the proof with a lemma.

Lemma 10.6. *Let $(\lambda_{nj})_{n=1,2,\dots,j=1,\dots,m_n}$ be a triangular family of asymptotically negligible, non-negative constants and $\lambda \in [0; \infty]$. Then applies*

$$\prod_{j=1}^{m_n} (1 - \lambda_{nj}) \xrightarrow{n \rightarrow \infty} e^{-\lambda} \quad \iff \quad \sum_{j=1}^{m_n} \lambda_{nj} \xrightarrow{n \rightarrow \infty} \lambda$$

Proof. First note that $\log(1 - x) = -x + \varepsilon(x)$ for $x > 0$ with $\varepsilon(x)/x \xrightarrow{x \rightarrow 0} 0$. Since $\sup_{j=1,\dots,m_n} \lambda_{nj} < 1$ for large n , the left statement is statement is equivalent to

$$-\lambda = \lim_{n \rightarrow \infty} \sum_{j=1}^{m_n} \log(1 - \lambda_{nj}) = - \lim_{n \rightarrow \infty} \sum_{j=1}^{m_n} \lambda_{nj} \left(1 - \frac{\varepsilon(\lambda_{nj})}{\lambda_{nj}}\right) = - \lim_{n \rightarrow \infty} \sum_{j=1}^{m_n} \lambda_{nj},$$

as

$$\sup_{j=1,\dots,m_n} \frac{\varepsilon(\lambda_{nj})}{\lambda_{nj}} \xrightarrow{n \rightarrow \infty} 0.$$

From this follows the statement. □

Proof of theorem 10.5. We denote by $\varphi_{n,j}$ the generating function of $X_{n,j}$. According to Remark ??.2, the convergence statement of the weak convergence in the theorem is equivalent to the pointwise convergence of $\prod_{j=1}^{m_n} \varphi_{n,j}(z) \xrightarrow{n \rightarrow \infty} e^{-\lambda(1-z)}$, since

$$\varphi_X(z) = \sum_{k=0}^{\infty} e^{-\lambda} \frac{\lambda^k}{k!} z^k = e^{-\lambda(1-z)}.$$

Because of Lemma 10.6 this is exactly true, if

$$A_n(z) := \sum_{j=1}^{m_n} (1 - \varphi_{n,j}(z)) \xrightarrow{n \rightarrow \infty} \lambda(1 - z), \quad (10.4)$$

since the family $(1 - \varphi_{n,j}(z))_{n=1,2,\dots,j=1,\dots,m_n}$ for each $z \in [0, 1]$ after (10.3) is asymptotically negligible. We decompose $A_n(z) = A_n^1(z) + A_n^2(z)$ with

$$A_n^1(z) = \sum_{k=1}^{\infty} (1 - z) \sum_{j=1}^{m_n} \mathbf{P}(X_{n,j} = k) = (1 - z) \sum_{j=1}^{m_n} \mathbf{P}(X_{n,j} > 0),$$

$$A_n^2(z) = \sum_{k=2}^{\infty} (z - z^k) \sum_{j=1}^{m_n} \mathbf{P}(X_{n,j} = k).$$

First, let it be established that $z(1 - z) \leq z - z^k \leq z$ for all $k = 2, 3, \dots$. This means that

$$z(1 - z) \sum_{j=1}^{m_n} \mathbf{P}(X_{n,j} > 1) \leq A_n^2(z) \leq z \sum_{j=1}^{m_n} \mathbf{P}(X_{n,j} > 1). \quad (10.5)$$

Let us now turn to the proof of the assertion.

' \Rightarrow ': Let (10.4) now apply. For $z = 0$ this means, since $\varphi_{nj}(0) = \mathbf{P}(X_{nj} = 0)$, that

$$\sum_{j=1}^{m_n} \mathbf{P}(X_{nj} > 0) = \sum_{j=1}^{m_n} (1 - \varphi_{nj}(0)) \xrightarrow{n \rightarrow \infty} \lambda,$$

therefore $A_n^1(z) \xrightarrow{n \rightarrow \infty} \lambda(1-z)$ for $z \in [0, 1]$. But then $A_n^2(z) \xrightarrow{n \rightarrow \infty} 0$ must apply to $z \in [0, 1]$. Because of (10.5) this means that 1. is valid. The statement 2. follows from this by subtraction. ' \Leftarrow ': So 1. and 2. apply. It is clear that $A_n^2(z) \xrightarrow{n \rightarrow \infty} 0$ because of (10.5). Then also applies $A_n^1(z) \xrightarrow{n \rightarrow \infty} (1-z)\lambda$ because of the condition, i.e. (10.4) is shown. \square

Example 10.7 (Convergence of geometric distributions against Poisson). *Let X_{nj} , $j = 1, \dots, n$, $n = 1, 2, \dots$ be geometrically distributed with parameter p_n (i.e. $\mathbf{P}(X_{nj} = k) = (1-p_n)^{k-1}p_n$, see example 2.2.4) and $Y_{nj} = X_{nj} - 1$. (Thus gives Y_{nj} the number of failures before the first success). We set $Y_n := \sum_{j=1}^n Y_{nj}$, which is as distributed as the number of failures before the n th success. If $Y \sim \text{Poi}(\lambda)$ and $(1-p_n) \cdot n \xrightarrow{n \rightarrow \infty} \lambda$, then $Y_n \xrightarrow{n \rightarrow \infty} Y$. Because the following applies*

$$\begin{aligned} \sum_{j=1}^n \mathbf{P}(Y_{nj} = 1) &= n(1-p_n)p_n \xrightarrow{n \rightarrow \infty} \lambda, \\ \sum_{j=1}^n \mathbf{P}(Y_{nj} > 1) &= n(1-p_n)^2 \xrightarrow{n \rightarrow \infty} 0 \end{aligned}$$

and theorem 10.5 gives the result.

10.2 The central limit theorem

The central limit theorem, Theorem 10.8, generalizes the theorem of deMoivre Laplace. The generalization consists of the fact that any sums of independent (not necessarily identically distributed) random variables converge weakly to a normally distributed random variable if they satisfy the *Lindeberg condition* (see 2. in Theorem 10.8) is fulfilled.

Theorem 10.8 (Central limit theorem of Lindeberg-Feller). *Let $(X_{nj})_{n=1,2,\dots,j=1,\dots,m_n}$ a family of random variables such that for $n = 1, 2, \dots$ the random variables X_{n1}, \dots, X_{nm_n} are independent. Let also be*

$$\sum_{j=1}^{m_n} \mathbf{E}[X_{nj}] \xrightarrow{n \rightarrow \infty} \mu, \quad \sum_{j=1}^{m_n} \mathbf{V}[X_{nj}] \xrightarrow{n \rightarrow \infty} \sigma^2$$

and $X \sim N(\mu, \sigma^2)$. Then the following statements are equivalent:

1. $\sum_{j=1}^{m_n} X_{nj} \xrightarrow{n \rightarrow \infty} X$ and $\sup_{j=1,\dots,m_n} \mathbf{V}[X_{nj}] \xrightarrow{n \rightarrow \infty} 0$,
2. $\sum_{j=1}^{m_n} \mathbf{E}[(X_{nj} - \mathbf{E}[X_{nj}])^2; |X_{nj} - \mathbf{E}[X_{nj}]| > \varepsilon] \xrightarrow{n \rightarrow \infty} 0$ for all $\varepsilon > 0$.

Before we prove the central limit theorem, we refer to the special case of identically distributed random variables, which was already discussed in the stochastics lecture.

Corollary 10.9 (Central limit theorem for identically distributed random variable). *Let X_1, X_2, \dots be independent and identically distributed with $\mathbf{E}[X_1] = \mu$, $\mathbf{V}[X_1] = \sigma^2 > 0$. Let $S_n := \sum_{k=1}^n X_k$ and $X \sim N(0, 1)$. Then applies*

$$\frac{S_n - n\mu}{\sqrt{n\sigma^2}} \xrightarrow{n \rightarrow \infty} X.$$

Proof. Let $m_n = n$ and $X_{nj} = \frac{X_j - \mu}{\sqrt{n\sigma^2}}$. Then the family $(X_{nj})_{n=1,2,\dots,j=1,\dots,n}$ fulfills the conditions of theorem 10.8 with $\mu = 0$, $\sigma^2 = 1$. Furthermore

$$\sum_{j=1}^n \mathbf{E}[X_{nj}^2; |X_{nj}| > \varepsilon] = \frac{1}{\sigma^2} \mathbf{E}[(X_1 - \mu)^2; |X_1 - \mu| > \varepsilon\sqrt{n\sigma^2}] \xrightarrow{n \rightarrow \infty} 0$$

due to majorized convergence. □

The Lindeberg condition is often not easy to verify. The stronger Lyapunoff condition is often simpler.

Remark 10.10 (Lyapunoff condition). *The family $(X_{nj})_{n=1,2,\dots,j=1,\dots,m_n}$ from Theorem 10.8 satisfies the Lyapunoff condition if for a $\delta > 0$*

$$\sum_{j=1}^{m_n} \mathbf{E}[|X_{nj} - \mathbf{E}[X_{nj}]|^{2+\delta}] \xrightarrow{n \rightarrow \infty} 0.$$

Under the conditions of Theorem 10.8, the Lyapunoff condition implies the Lindeberg condition. To see this, let without restriction $\mathbf{E}[X_{nj}] = 0$. The following applies for all $\varepsilon > 0$

$$x^2 1_{|x|>\varepsilon} \leq \frac{|x|^{2+\delta}}{\varepsilon^\delta} 1_{|x|>\varepsilon} \leq \frac{|x|^{2+\delta}}{\varepsilon^\delta}.$$

If the Lyapunoff condition applies, the Lindeberg condition follows from

$$\sum_{j=1}^{m_n} \mathbf{E}[X_{nj}^2; |X_{nj}| > \varepsilon] \leq \frac{1}{\varepsilon^\delta} \sum_{j=1}^{m_n} \mathbf{E}[|X_{nj}|^{2+\delta}] \xrightarrow{n \rightarrow \infty} 0.$$

The proof of theorem 10.8 is based on the clever use of the characteristic functions of the random variable random variable X_{nj} and Taylor approximations. We prepare the proof of the theorem with two lemmas.

Lemma 10.11 (estimate). *For complex numbers $z_1, \dots, z_n, z'_1, \dots, z'_n$ with $|z_i| \leq 1, |z'_i| \leq 1$ for $i = 1, \dots, n$ applies*

$$\left| \prod_{k=1}^n z_k - \prod_{k=1}^n z'_k \right| \leq \sum_{k=1}^n |z_k - z'_k|. \quad (10.6)$$

Proof. For $n = 1$ the equation is obviously correct. Applies (10.6) is valid for an n , then

$$\begin{aligned} \left| \prod_{k=1}^{n+1} z_k - \prod_{k=1}^{n+1} z'_k \right| &\leq \left| z_{n+1} \left(\prod_{k=1}^n z_k - \prod_{k=1}^n z'_k \right) \right| + \left| (z_{n+1} - z'_{n+1}) \prod_{k=1}^n z'_k \right| \\ &\leq \sum_{k=1}^n |z_k - z'_k| + |z_{n+1} - z'_{n+1}|. \end{aligned}$$

From this follows the assertion. □

Lemma 10.12 (Taylor approximation of the exponential function). *Let $t \in \mathbb{C}$ and $n \in \mathbb{Z}_+$. Then applies*

$$\left| e^{it} - \sum_{k=0}^n \frac{(it)^k}{k!} \right| \leq \frac{2|t|^n}{n!} \wedge \frac{|t|^{n+1}}{(n+1)!}. \quad (10.7)$$

Proof. Denote $h_n(t)$ the difference on the left-hand side. For $n = 0$ (10.7) follows from

$$|h_0(t)| = \left| \int_0^t e^{is} ds \right| \leq \int_0^t |e^{is}| ds = |t|$$

and

$$|h_0(t)| \leq |e^{it}| + 1 = 2.$$

In general, the following applies to $t \in \mathbb{R}, n \in \mathbb{N}$

$$\left| \int_0^t h_n(s) ds \right| = \left| -i(e^{it} - 1) + i \sum_{k=0}^n \frac{(it)^{k+1}}{(k+1)!} \right| = \left| ie^{it} - i \sum_{k=0}^{n+1} \frac{(it)^k}{k!} \right| = |h_{n+1}(t)|,$$

from which (10.7) follows by induction. \square

Remark 10.13 (notation). *In the following proof, we will prove for functions a and b exactly when $a \leq b$ if there is a constant C with $a \leq Cb$.*

Proof of theorem 10.8. O.E. let $\mathbf{E}[X_{nj}] = \mu = 0$ and $\sigma^2 = 1$; otherwise we replace X_{nj} by $\frac{X_{nj} - \mathbf{E}[X_{nj}]}{\sqrt{\sigma^2}}$. Let $\sigma_{nj}^2 := \mathbf{V}[X_{nj}]$ and $\sigma_n^2 := \sum_{j=1}^{m_n} \sigma_{nj}^2 \xrightarrow{n \rightarrow \infty} 1$. Denote also ψ_{nj} the characteristic function of X_{nj} .

2. \Rightarrow 1. Since for every $\varepsilon > 0$

$$\sup_{j=1, \dots, m_n} \sigma_{nj}^2 \leq \varepsilon^2 + \sup_{j=1, \dots, m_n} \mathbf{E}[X_{nj}^2; |X_{nj}| > \varepsilon] \leq \varepsilon^2 + \sum_{j=1}^{m_n} \mathbf{E}[X_{nj}^2; |X_{nj}| > \varepsilon] \xrightarrow{n \rightarrow \infty} \varepsilon^2, \quad (10.8)$$

the second part of 1. is already shown.

Let $(Z_{nj})_{n=1,2,\dots,j=1,\dots,m_n}$ be independent random variables with $Z_{nj} \sim N(0, \sigma_{nj}^2)$. This means that $Z_n = \sum_{j=1}^{m_n} Z_{nj} \sim N(0, \sigma_n^2)$. In particular, the following applies thus $Z_n \xrightarrow{n \rightarrow \infty} X$, which can be derived directly from the form of the characteristic functions of the normal distribution, Example 6.13.3 can be read off. Let $\tilde{\psi}_{nj}$ be the characteristic function of Z_{nj} . Then it suffices to show, see theorem 9.33, that

$$\prod_{j=1}^{n_j} \psi_{nj}(t) - \prod_{j=1}^{m_n} \tilde{\psi}_{nj}(t) \xrightarrow{n \rightarrow \infty} 0 \quad (10.9)$$

for all t . Using Lemma 10.11 and Lemma 10.12 we write

$$\begin{aligned} \left| \prod_{j=1}^{m_n} \psi_{nj}(t) - \prod_{j=1}^{m_n} \tilde{\psi}_{nj}(t) \right| &\leq \sum_{j=1}^{m_n} |\psi_{nj}(t) - \tilde{\psi}_{nj}(t)| \\ &\leq \sum_{j=1}^{m_n} \left| \psi_{nj}(t) - 1 + \frac{1}{2}t^2\sigma_{nj}^2 \right| + \sum_{j=1}^{m_n} \left| \tilde{\psi}_{nj}(t) - 1 + \frac{1}{2}t^2\sigma_{nj}^2 \right| \\ &\lesssim 2 \sum_{j=1}^{m_n} \mathbf{E}[X_{nj}^2(1 \wedge |X_{nj}|)] + \sum_{j=1}^{m_n} \left| e^{-\frac{1}{2}\sigma_{nj}^2 t^2} - 1 + \frac{1}{2}t^2\sigma_{nj}^2 \right|. \end{aligned}$$

Further is

$$\sum_{j=1}^{m_n} \mathbf{E}[X_{nj}^2(1 \wedge |X_{nj}|)] \leq \varepsilon \sum_{j=1}^{m_n} \sigma_{nj}^2 + \sum_{j=1}^{m_n} \mathbf{E}[X_{nj}^2; |X_{nj}| > \varepsilon] \xrightarrow{n \rightarrow \infty} \varepsilon$$

and

$$\sum_{j=1}^{m_n} |e^{-\frac{1}{2}\sigma_{nj}^2 t^2} - 1 + \frac{1}{2}t^2 \sigma_{nj}^2| \lesssim \sum_{j=1}^{m_n} \sigma_{nj}^4 \leq \sigma_n^2 \sup_{j=1, \dots, m_n} \sigma_{nj}^2 \xrightarrow{n \rightarrow \infty} 0$$

because of (10.8). This means (10.9) already proven.

1. \Rightarrow 2. According to the second part of 1. for each $\varepsilon > 0$ with the Chebyshev inequality

$$\sup_{j=1, \dots, m_n} \mathbf{P}[|X_{nj}| > \varepsilon] \leq \sup_{j=1, \dots, m_n} \frac{\sigma_{nj}^2}{\varepsilon^2} \xrightarrow{n \rightarrow \infty} 0. \quad (10.10)$$

With Lemma 10.12 applies

$$\sup_{j=1, \dots, m_n} |\psi_{nj}(t) - 1| \leq \sup_{j=1, \dots, m_n} \mathbf{E}[2 \wedge |t \cdot X_{nj}|] \leq 2 \sup_{j=1, \dots, m_n} \mathbf{P}[|X_{nj}| > \varepsilon] + \varepsilon |t| \xrightarrow{n \rightarrow \infty} \varepsilon |t|.$$

In particular, $\sum_{j=1}^{m_n} \log \psi_{nj}(t)$ is defined for every t if n is large enough. From the validity of 1. follows

$$\sum_{j=1}^{m_n} \log \psi_{nj}(t) \xrightarrow{n \rightarrow \infty} -\frac{t^2}{2}. \quad (10.11)$$

Furthermore, because $\psi'_{nj}(0) = i\mathbf{E}[X_{nj}] = 0$, $\psi''_{nj}(0) = -\mathbf{V}[X_{nj}] = -\sigma_{nj}^2$ with the help of a Taylor expansion of ψ_{nj} around 0

$$|\psi_{nj}(t) - 1| \lesssim \sigma_{nj}^2 |t|^2$$

and

$$\begin{aligned} \left| \sum_{j=1}^{m_n} \log \psi_{nj}(t) - \sum_{j=1}^{m_n} (\psi_{nj}(t) - 1) \right| &\lesssim \sum_{j=1}^{m_n} |\psi_{nj}(t) - 1|^2 \\ &\lesssim \sum_{j=1}^{m_n} (\sigma_{nj}^2)^2 |t|^4 \lesssim |t|^4 \sup_{j=1, \dots, m_n} \sigma_{nj}^2 \xrightarrow{n \rightarrow \infty} 0. \end{aligned} \quad (10.12)$$

Since the convergence of an imaginary series follows from the convergence of its real part follows, we deduce from (10.11) and (10.12) because $\operatorname{Re}(\psi_{nj}(t)) = \mathbf{E}[\cos(tX_{nj})]$

$$\sum_{j=1}^{m_n} \mathbf{E}[\cos(tX_{nj}) - 1] \xrightarrow{n \rightarrow \infty} -\frac{t^2}{2}$$

For $\varepsilon > 0$ is now because of $0 \leq 1 - \cos(\theta) \leq \frac{\theta^2}{2}$

$$\begin{aligned}
0 \leq \limsup_{n \rightarrow \infty} \sum_{j=1}^{m_n} \mathbf{E}[X_{nj}^2; |X_{nj}| > \varepsilon] &= \limsup_{n \rightarrow \infty} 1 - \sum_{j=1}^{m_n} \mathbf{E}[X_{nj}^2; |X_{nj}| \leq \varepsilon] \\
&\leq \limsup_{n \rightarrow \infty} 1 - \frac{2}{t^2} \sum_{j=1}^{m_n} \mathbf{E}[1 - \cos(tX_{nj}); |X_{nj}| \leq \varepsilon] \\
&= \limsup_{n \rightarrow \infty} \frac{2}{t^2} \sum_{j=1}^{m_n} \mathbf{E}[1 - \cos(tX_{nj}); |X_{nj}| > \varepsilon] \quad (10.13) \\
&\leq \limsup_{n \rightarrow \infty} \frac{2}{t^2} \sum_{j=1}^{m_n} \mathbf{P}[|X_{nj}| > \varepsilon] \\
&\leq \frac{2}{\varepsilon^2 t^2} \limsup_{n \rightarrow \infty} \sum_{j=1}^{m_n} \sigma_{nj} = \frac{2}{\varepsilon^2 t^2}.
\end{aligned}$$

Since $t, \varepsilon > 0$ were arbitrary, 2. is shown, if in the the last inequality chain $t \rightarrow \infty$ is considered. \square

10.3 Multidimensional limit sets

So far, we have only considered weak limit theorems for the case of \mathbb{R} -valued random variables. We now generalize this to \mathbb{R}^d -valued random variables. In particular, we give a variant of the multidimensional central limit theorem.

Definition 10.14 (Multidimensional normal distribution). *Let $\mu \in \mathbb{R}^d$ and $C \in \mathbb{R}^{d \times d}$ be a strictly positive definite symmetric matrix.^{4,5} The d -dimensional normal distribution with expected value μ and covariance matrix C is the probability measure $N_{\mu, C}$ on \mathbb{R}^d with density*

$$f_{\mu, C}(x) = \frac{1}{\sqrt{(2\pi)^d \det(C)}} \exp\left(-\frac{1}{2}(x - \mu)C^{-1}(x - \mu)^\top\right).$$

Proposition 10.15 (Properties of the multidimensional normal distribution). *Let $\mu \in \mathbb{R}^d$, $C = AA^\top \in \mathbb{R}^{d \times d}$ a strictly positive definite symmetric matrix and I the d -dimensional unit matrix. They are equivalent:*

1. $X \sim N_{\mu, C}$
2. $tX^\top \sim N_{t\mu^\top, tCt^\top}$ for each $t \in \mathbb{R}^d$
3. $\psi_X(t) = e^{it\mu^\top} e^{-\frac{1}{2}tCt^\top}$ for each $t \in \mathbb{R}^d$.

In each of these cases

4. $X \stackrel{d}{=} AY + \mu$ for $Y \sim N_{0, I}$

⁴We denote here row vectors with here denote row vectors by x and column vectors by x^\top .

⁵Strict positive definite means $xCx^\top > 0$ for all $x \in \mathbb{R}^d$. From linear algebra it is known that for a strictly positive definite matrix C there is always an invertible matrix A with $C = AA^\top$

5. $\mathbf{E}[X_i] = \mu_i$ for $i = 1, \dots, d$

6. $\mathbf{COV}[X_i, X_j] = C_{ij}$ for $i, j = 1, \dots, d$

Proof. First, let $X \sim N_{\mu, C}$. We first show 4.-6. The property 4. is an application of the transformation theorem. For $B \in \mathcal{B}(\mathbb{R}^d)$ and $T : y \mapsto Ay^\top + \mu^\top$ is

$$\begin{aligned} N_{0, I}(T^{-1}(B)) &= \frac{1}{\sqrt{(2\pi)^d}} \int_{T^{-1}(B)} e^{-\frac{1}{2}yy^\top} dy \\ &\stackrel{y=A^{-1}(x-\mu)}{=} \frac{1}{\sqrt{(2\pi)^d}} \frac{1}{\det A} \int_B \exp\left(-\frac{1}{2}(x-\mu)(A^\top)^{-1}A^{-1}(x-\mu)^\top\right) dx \\ &= \frac{1}{\sqrt{(2\pi)^d \det C}} \int_B \exp\left(-\frac{1}{2}(x-\mu)C^{-1}(x-\mu)^\top\right) dx \\ &= N_{\mu, C}(B). \end{aligned}$$

5. follows from 4. with

$$\mathbf{E}[X_i] = \mathbf{E}[\pi_i(Ay + \mu)] = \pi_i\mu = \mu_i,$$

where π_i is the projection onto the i -th coordinate.

6. also follows from 4. with

$$\begin{aligned} \mathbf{COV}[X_i, X_j] &= \mathbf{E}[(\pi_i AY^\top)(\pi_j AY^\top)] = \mathbf{E}[(A_i \cdot Y^\top)(A_j \cdot Y^\top)] = \mathbf{E}[A_i \cdot Y^\top Y A_j^\top] \\ &= A_i \cdot A_j^\top = (AA^\top)_{ij} = C_{ij}. \end{aligned}$$

We now come to the equivalence of 1.-3.: '1. \Rightarrow 2.': Since $X \stackrel{d}{=} AY^\top + \mu^\top$ as in 4. $tX^\top = tAY^\top + t\mu^\top$ as a linear combination of (one-dimensional) normal distributions is normally distributed again. The expected value is obviously $t\mu^\top$ and the variance

$$\mathbf{V}[tX^\top] = \mathbf{E}[(tAY^\top)^2] = \mathbf{E}[tAY^\top Y A^\top t^\top] = tAA^\top t^\top = tCt^\top.$$

'2. \Rightarrow 3.': Since $tX^\top \sim N_{t\mu^\top, tCt^\top}$, the statement follows from example 6.13.3.

'3. \Rightarrow 1.': This follows from Proposition 9.25. \square

Remark 10.16 (special cases). 1. If C in Definition 10.14 is positive, but not strictly positive definite strictly positive definite (i.e. there is $x \in \mathbb{R}^d$, $x \neq 0$ with $xCx = 0$), one cannot determine $N_{\mu, C}$ by specifying the density as in the definition above. In this case $N_{\mu, C}$ is defined by specifying the characteristic function, i.e. function, i.e. $N_{\mu, C}$ is the uniquely determined distribution on \mathbb{R}^d with $\psi_{N_{\mu, C}}(t) = e^{it\mu} e^{-\frac{1}{2}tCt^\top}$.

2. If $Y \sim N_{0, I}$ and A is an orthogonal matrix, then is also $X := AY \sim N_{0, I}$. This follows from Proposition 10.15, if you write $I = AA^\top$ and use 4. is used.

Proposition 10.17 (Cramér-Wold Device). If X, X_1, X_2, \dots are random variables with values in \mathbb{R}^d . Then $X_n \xrightarrow{n \rightarrow \infty} X$ applies if and only if $tX_n \xrightarrow{n \rightarrow \infty} tX$ for all $t \in \mathbb{R}^d$ (where $(t, x) \mapsto tx$ is the scalar product in \mathbb{R}^d).

Proof. ' \Rightarrow ': Let $t \in \mathbb{R}^d$ and $f \in C_b(\mathbb{R})$. Then $f(t \cdot) \in C_b(\mathbb{R}^d)$. This means that $\mathbf{E}[f(tX_n)] \xrightarrow{n \rightarrow \infty} \mathbf{E}[f(tX)]$, i.e. $tX_n \xrightarrow{n \rightarrow \infty} tX$.

' \Leftarrow ': Let π_i be the projection onto the i th coordinate. Since $(\pi_i X_n)_{n=1,2,\dots}$ according to Corollary 9.18 is tight for all i , you can see that $(X_n)_{n=1,2,\dots}$ is tight. Since $\{x \mapsto e^{itx} : t \in \mathbb{R}^d\}$ is a separating class of functions, it follows that class of functions, the assertion follows from $\mathbf{E}[e^{itX_n}] \xrightarrow{n \rightarrow \infty} \mathbf{E}[e^{itX}]$ for all $t \in \mathbb{R}^d$ and Proposition 9.27. \square

Theorem 10.18 (Multidimensional central limit theorem). *Let X_1, X_2, \dots be independent, identical distributed random variables with values in \mathbb{R}^d with $\mathbf{E}[X_n] = \mu \in \mathbb{R}^d$ and $\mathbf{COV}[X_{n,i}, X_{n,j}] = C_{ij}$ for $i, j = 1, \dots, d$ and $S_n = \sum_{i=1}^n X_i$. If $X \sim N_{0,C}$, then the following applies*

$$\frac{S_n - n\mu}{\sqrt{n}} \xrightarrow{n \rightarrow \infty} X.$$

Proof. We apply the one-dimensional central limit theorem, Corollary 10.9, to the independent, identically distributed random variables tX_1, tX_2, \dots . This provides

$$t \frac{S_n - n\mu}{\sqrt{n}} \xrightarrow{n \rightarrow \infty} tX.$$

Since t was arbitrary, the statement follows from Proposition 10.17. \square

11 The conditional expectation

Let $(\Omega, \mathcal{A}, \mathbf{P})$ be a probability space. We write $\mathcal{L}^1 := \mathcal{L}^1(\mathbf{P})$ for the set of all real random variables whose expected value exists. In this chapter we again use the notation $\mathbb{E}[\cdot]$ for the integral with respect to the probability measure \mathbf{P} , as well as $\mathcal{L}^p := \mathcal{L}^p(\mathbf{P})$.

11.1 motivation

Define as in elementary stochastics for $A, G \in \mathcal{A}$ and $\mathbf{P}(G) > 0$

$$\mathbf{P}(A|G) := \frac{\mathbf{P}(A \cap G)}{\mathbf{P}(G)}$$

and analogously the *conditional expectation*

$$\mathbf{E}[X|G] := \frac{\mathbf{E}[X; G]}{\mathbf{P}(G)}.$$

Then $\mathbf{P}(A|G) = \mathbf{E}[1_A|G]$. This relationship means that conditional expectations can be used to calculate conditional probabilities. In particular, the notion of conditional expectation is more general than the notion of conditional probability.

In this chapter, we will use the conditional expectation $\mathbf{E}[X|\mathcal{G}]$ for a random variable X and a σ algebra $\mathcal{G} \subseteq \mathcal{F}$ learn. Here, $\mathbf{E}[X|\mathcal{G}]$ is a \mathcal{G} -measurable random variable. As a simple example $\{G_1, G_2, \dots\} \subseteq \mathcal{F}$ is a partition of Ω with $\mathbf{P}(G_i) > 0$ for $i = 1, 2, \dots$ and \mathcal{G} the generated σ algebra. Then we set for $X \in \mathcal{L}^1$

$$\mathbf{E}[X|\mathcal{G}](\omega) := \sum_{i=1}^{\infty} \mathbf{E}[X|G_i] \cdot 1_{G_i}(\omega). \quad (11.1)$$

The following therefore applies: for $\omega \in G_i$, the random variable $\mathbf{E}[X|\mathcal{G}]$ is given by $\mathbf{E}[X|\mathcal{G}](\omega) = \mathbf{E}[X|G_i] = \mathbf{E}[X; G_i]/\mathbf{P}(G_i)$. In particular it is constant on G_i , $i = 1, 2, \dots$. In other words,

$\mathbf{E}[X|\mathcal{G}]$ is measurable with respect to \mathcal{G} . The following also applies to $J \subseteq \mathbb{N}$ and $A = \bigcup_{j \in J} G_j \in \mathcal{G}$

$$\begin{aligned} \mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A] &= \mathbf{E}\left[\sum_{i=1}^{\infty} \mathbf{E}[X|G_i] 1_{G_i} 1_A\right] \\ &= \sum_{j \in J} \mathbf{E}[\mathbf{E}[X|G_j] 1_{G_j}] \\ &= \sum_{j \in J} \mathbf{E}[X|G_j] \cdot \mathbf{P}(G_j) \\ &= \mathbf{E}[X; A]. \end{aligned} \tag{11.2}$$

In particular, with $J = \mathbb{N}$ therefore $\mathbf{E}[\mathbf{E}[X|\mathcal{F}]] = \mathbf{E}[X]$. The definition of the conditional expectation (11.1) can be defined with the help of the property (11.2) to any σ -algebras $\mathcal{G} \subseteq \mathcal{F}$ can be generalized.

Example 11.1 (binomial distribution with random probability of success). *Let X be uniformly distributed on $[0, 1]$, i.e. the distribution of X has density $1_{[0;1]}$. Given $X = x$ let Y_1, \dots, Y_n be a sequence of Bernoulli distributed random variables with probability of success x . Therefore, $Y = Y_1 + \dots + Y_n$ is binomially distributed with n and x , i.e. Y counts the number of successes in n independent experiments with probability of success x . Intuitively, it is clear that the*

$$\mathbf{P}(Y = k|X) = \binom{n}{k} X^k (1 - X)^{n-k}$$

should mean. However, this has not yet been defined, since $\mathbf{P}(X = x) = 0$ applies. However, it is worth noting that the right side is a $\sigma(X)$ -measurable random variable.

11.2 definition and properties

We now formally define the conditional expectation $\mathbf{E}[X|\mathcal{G}]$ for $\mathcal{G} \subseteq \mathcal{F}$. As mentioned above, this is this is a \mathcal{G} -measurable random variable whose expectations are as in (11.2) match those of X .

Theorem 11.2 (Existence and properties of the conditional expectation). *Let $\mathcal{G} \subseteq \mathcal{F}$ be a σ -algebra. Then there is there is an almost surely unique linear operator $\mathbf{E}[\cdot|\mathcal{G}] : \mathcal{L}^1 \rightarrow \mathcal{L}^1$ such that $\mathbf{E}[X|\mathcal{G}]$ for all $X \in \mathcal{L}^1$ a \mathcal{G} -measurable random variable with*

1. $\mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A] = \mathbf{E}[X; A]$ for all $A \in \mathcal{G}$.

Further applies

2. $\mathbf{E}[X|\mathcal{G}] \geq 0$ if $X \geq 0$.
3. $\mathbf{E}[|\mathbf{E}[X|\mathcal{G}]|] \leq \mathbf{E}[|X|]$.
4. If $0 \leq X_n \uparrow X$ for $n \rightarrow \infty$, then also $\mathbf{E}[X_n|\mathcal{G}] \uparrow \mathbf{E}[X|\mathcal{G}]$ in \mathcal{L}^1 if all expectations exist.
5. If X is a \mathcal{G} -measurable function, then $\mathbf{E}[XY|\mathcal{G}] = X\mathbf{E}[Y|\mathcal{G}]$ if all expectations exist.
6. $\mathbf{E}[X\mathbf{E}[Y|\mathcal{G}]] = \mathbf{E}[\mathbf{E}[X|\mathcal{G}]Y] = \mathbf{E}[\mathbf{E}[X|\mathcal{G}]\mathbf{E}[Y|\mathcal{G}]]$ if all expectations exist. expectations exist.

7. If $\mathcal{H} \subseteq \mathcal{G}$, then $\mathbf{E}[\mathbf{E}[X|\mathcal{G}]|\mathcal{H}] = \mathbf{E}[X|\mathcal{H}]$.

8. If X is independent of \mathcal{G} , then $\mathbf{E}[X|\mathcal{G}] = \mathbf{E}[X]$.

Proof. 1. in the case $X \in \mathcal{L}^2$: Let M be the closed linear subspace of \mathcal{L}^2 , which consists of all functions which, except for a zero set, correspond to a \mathcal{G} -measurable function. According to Proposition 4.10 there are now almost certainly unique functions $Y \in M, Z \perp M$ with $X = Y + Z$. We define $\mathbf{E}[X|\mathcal{G}] := Y$. This means that $X - \mathbf{E}[X|\mathcal{G}] \perp M$, i.e. $\mathbf{E}[X - \mathbf{E}[X|\mathcal{G}]; A] = 0$ for $A \in \mathcal{G}$, from which 1. for $X \in \mathcal{L}^2$ follows.

3. in the case $X \in \mathcal{L}^2$: Choose $A := \{\mathbf{E}[X|\mathcal{G}] \geq 0\}$. According to 1. then applies

$$\mathbf{E}[|\mathbf{E}[X|\mathcal{G}]|] = \mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A] - \mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A^c] = \mathbf{E}[X; A] - \mathbf{E}[X; A^c] \leq \mathbf{E}[|X|].$$

1. in the case $X \in \mathcal{L}^1$: If $X \in \mathcal{L}^1 \supset \mathcal{L}^2$, then choose $X_1, X_2, \dots \in \mathcal{L}^2$ with $\|X_n - X\|_1 \xrightarrow{n \rightarrow \infty} 0$ (such that $|X_n| := |X| \wedge n$), and define $\mathbf{E}[X|\mathcal{G}] := \lim_{n \rightarrow \infty} \mathbf{E}[X_n|\mathcal{G}]$. This limit value exists in \mathcal{L}^1 , since because of 3.

$$\mathbf{E}[|\mathbf{E}[X_n|\mathcal{G}] - \mathbf{E}[X_m|\mathcal{G}]|] = \mathbf{E}[|\mathbf{E}[X_n - X_m|\mathcal{G}]|] \leq \mathbf{E}[|X_n - X_m|] \xrightarrow{n, m \rightarrow \infty} 0$$

the sequence $(\mathbf{E}[X_n|\mathcal{G}])_{n=1,2,\dots}$ is a Cauchy sequence and \mathcal{L}^1 is complete. Furthermore, this means that $\|\mathbf{E}[X_n|\mathcal{G}] - \mathbf{E}[X|\mathcal{G}]\|_1 \xrightarrow{n \rightarrow \infty} 0$. Furthermore, for $A \in \mathcal{G}$

$$\begin{aligned} |\mathbf{E}[X - \mathbf{E}[X|\mathcal{G}]; A]| &\leq \mathbf{E}[|X1_A - X_n1_A|] \\ &\quad + |\mathbf{E}[X_n - \mathbf{E}[X_n|\mathcal{G}]; A]| \\ &\quad + \mathbf{E}[|\mathbf{E}[X_n|\mathcal{G}]1_A - \mathbf{E}[X|\mathcal{G}]1_A|] \\ &\xrightarrow{n \rightarrow \infty} 0 \end{aligned}$$

due to majorized convergence and 1. follows in the case $X \in \mathcal{L}^1$.

3. in the case $X \in \mathcal{L}^1$. Here, too, you can see through an approximation argument if $X_1, X_2, \dots \in \mathcal{L}^2$ with $X_n \xrightarrow{n \rightarrow \infty} \mathcal{L}^1 X$

$$\mathbf{E}[|\mathbf{E}[X|\mathcal{G}]|] = \lim_{n \rightarrow \infty} \mathbf{E}[|\mathbf{E}[X_n|\mathcal{G}]|] \leq \lim_{n \rightarrow \infty} \mathbf{E}[|X_n|] = \mathbf{E}[|X|],$$

since, due to the inverse triangle inequality, approximately

$$\mathbf{E}[|\mathbf{E}[X_n|\mathcal{G}]| - |\mathbf{E}[X|\mathcal{G}]|] \leq \mathbf{E}[|\mathbf{E}[X|\mathcal{G}] - \mathbf{E}[X_n|\mathcal{G}]|] \xrightarrow{n \rightarrow \infty} 0.$$

2. set $A = \{\mathbf{E}[X|\mathcal{G}] \leq 0\}$ and thus

$$0 \geq \mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A] = \mathbf{E}[X; A] \geq 0,$$

thus because of $\mathbf{E}[X|\mathcal{G}]1_A \leq 0$ also $\mathbf{E}[X|\mathcal{G}]1_A = 0$ is almost certain.

4 Due to monotonic convergence $\|X_n - X\|_1 \xrightarrow{n \rightarrow \infty} 0$, i.e. after 3.

$$\mathbf{E}[|\mathbf{E}[X_n|\mathcal{G}] - \mathbf{E}[X|\mathcal{G}]|] = \mathbf{E}[|\mathbf{E}[X_n - X|\mathcal{G}]|] \leq \mathbf{E}[|X_n - X|] \xrightarrow{n \rightarrow \infty} 0.$$

6. in the case $X, Y \in \mathcal{L}^2$. According to the definition of the conditional expectation is $\mathbf{E}[X|\mathcal{G}], \mathbf{E}[Y|\mathcal{G}] \in M$ if M is the linear subspace of \mathcal{L}^2 which contains functions which, apart from a zero set with a \mathcal{G} -measurable function. Furthermore $X - \mathbf{E}[X|\mathcal{G}] \perp M$. Thus

$$\mathbf{E}[(X - \mathbf{E}[X|\mathcal{G}])\mathbf{E}[Y|\mathcal{G}]] = 0.$$

6. in the case $X, Y \in \mathcal{L}^1$. Wähle $X_1, Y_1, X_2, Y_2, \dots \in \mathcal{L}^2$ with $X_n \uparrow X, Y_n \uparrow Y$. Because of 4. and majorized convergence, if all expectations exist,

$$\mathbf{E}[(X - \mathbf{E}[X|\mathcal{G}])\mathbf{E}[Y|\mathcal{G}]] = \lim_{n \rightarrow \infty} \mathbf{E}[(X_n - \mathbf{E}[X_n|\mathcal{G}])\mathbf{E}[Y_n|\mathcal{G}]] = 0.$$

5. because of 1. is $\mathbf{E}[X|\mathcal{G}]1_A = X1_A$ for $A \in \mathcal{G}$ is almost certain. This means that

$$\mathbf{E}[XY; A] = \mathbf{E}[X\mathbf{E}[Y|\mathcal{G}]; A]$$

after 6. from this follows after 1. already $\mathbf{E}[XY|\mathcal{G}] = X\mathbf{E}[Y|\mathcal{G}]$.

Since $\mathcal{H} \subseteq \mathcal{G}$, for $A \in \mathcal{H}$

$$\mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A] = \mathbf{E}[X; A] = \mathbf{E}[\mathbf{E}[X|\mathcal{H}]; A]$$

after 1. From here follows but $\mathbf{E}[\mathbf{E}[X|\mathcal{G}]|\mathcal{H}] = \mathbf{E}[X|\mathcal{H}]$.

8 Certainly $\mathbf{E}[X]$ is measurable with respect to \mathcal{G} . For $A \in \mathcal{G}$ is also

$$\mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A] = \mathbf{E}[X; A] = \mathbf{E}[X]\mathbf{E}[1_A] = \mathbf{E}[\mathbf{E}[X]; A]$$

and thus $\mathbf{E}[X|\mathcal{G}] = \mathbf{E}[X]$. □

Remark 11.3 (Interpretation and alternative proof). 1. Let $X \in \mathcal{L}^2$. As the proof of 1. in Theorem 11.2 shows, $X - \mathbf{E}[X|\mathcal{G}]$ is perpendicular to the linear subspace of all subspace of all \mathcal{G} -measurable functions. In particular $\mathbf{E}[X|\mathcal{G}]$ is the \mathcal{G} -measurable random variable that (in terms of the \mathcal{L}^2 norm) is closest to the random variable X comes closest. Therefore, we can say that $\mathbf{E}[X|\mathcal{G}]$ is the best estimate of X if information from the σ algebra \mathcal{G} is available. are available.

2. The almost certainly unambiguous existence of the conditional expectation with the property 1. in Theorem 11.2 can be proved differently than above with the help of the theorem of Radon-Nikodým (Corollary 4.17):

Let $X \geq 0$ first. Set $\tilde{\mathbf{P}} := \mathbf{P}|_{\mathcal{G}}$, the restriction of \mathbf{P} to \mathcal{G} , and $\mu(\cdot) := \tilde{\mathbf{E}}[X; \cdot]$ a finite measure. Then obviously $\mu \ll \tilde{\mathbf{P}}$ applies. The theorem of Radon-Nikodým ensures that μ is a density with respect to $\tilde{\mathbf{P}}$, i.e. there is a \mathcal{G} -measurable random variable Z with

$$\mathbf{E}[X; A] = \tilde{\mathbf{E}}[X; A] = \mu(A) = \tilde{\mathbf{E}}[Z; A] = \mathbf{E}[Z; A]$$

for all $A \in \mathcal{G}$. Thus Z fulfills the properties of 1. from theorem 11.2. The general case (i.e. X can also take can also assume negative values) then follows with the decomposition $X = X^+ - X^-$.

To prove the (almost certain) uniqueness of the conditional expectation expectation, let Z' be another \mathcal{G} -measurable random variable with random variable with $\mathbf{E}[Z'; A] = \mathbf{E}[X; A]$ for all $A \in \mathcal{G}$. Then $B := \{Z' - \mathbf{E}[X|\mathcal{G}] > 0\} \in \mathcal{G}$ and $\mathbf{E}[\mathbf{E}[X|\mathcal{G}] - Z'; B] = \mathbf{E}[X - X; B] = 0$ and likewise $\mathbf{E}[\mathbf{E}[X|\mathcal{G}] - Z'; B^c] = 0$. This therefore means $Z' = \mathbf{E}[X|\mathcal{G}]$ is almost certain.

Proposition 11.4 (Jensen's inequality for conditional expectations). Let I be an open interval, $\mathcal{G} \subseteq \mathcal{A}$ and $X \in \mathcal{L}^1$ with values in I and $\varphi : I \rightarrow \mathbb{R}$ is convex. Then applies

$$\mathbf{E}[\varphi(X)|\mathcal{G}] \geq \varphi(\mathbf{E}[X|\mathcal{G}]).$$

Proof. The proof is analogous to that of Jensen's inequality in the unconditional case, Proposition 6.6: Since I is open, $\mathbf{E}[X|\mathcal{G}] \in I$ is almost certain. We recall the definition of λ in (6.4). Further, as in (6.5) for $x \in I$

$$\varphi(x) \geq \varphi(\mathbf{E}[X|\mathcal{G}]) + \lambda(\mathbf{E}[X|\mathcal{G}]) (x - \mathbf{E}[X|\mathcal{G}])$$

and thus

$$\begin{aligned} \mathbf{E}[\varphi(X)|\mathcal{G}] &\geq \mathbf{E}[\varphi(\mathbf{E}[X|\mathcal{G}])|\mathcal{G}] + \mathbf{E}[\lambda(\mathbf{E}[X|\mathcal{G}]) \cdot (X - \mathbf{E}[X|\mathcal{G}])|\mathcal{G}] \\ &= \varphi(\mathbf{E}[X|\mathcal{G}]). \end{aligned} \quad \square$$

Lemma 11.5 (Equal integrability and conditional expectation). *Let $X \in \mathcal{L}^1$. Then the family $(\mathbf{E}[X|\mathcal{G}])_{\mathcal{G} \subseteq \mathcal{A}}$ is integrable to the same degree.*

Proof. Since $\{X\}$ is integrable to the same degree, according to lemma 7.9 there is a monotonically increasing convex function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\frac{\varphi(x)}{x} \xrightarrow{x \rightarrow \infty} \infty$ and $\mathbf{E}[\varphi(|X|)] < \infty$. With theorem 11.2.3 follows

$$\sup_{\mathcal{F} \subseteq \mathcal{A}} \mathbf{E}[\varphi(|\mathbf{E}[X|\mathcal{F}]|)] \leq \mathbf{E}[\varphi(|X|)] < \infty.$$

This means that $\{\mathbf{E}[X|\mathcal{F}] : \mathcal{F} \subseteq \mathcal{A} \text{ } \sigma\text{-algebra}\}$ equally integrable, again according to lemma 7.9. \square

Theorem 11.6 (Majorized and monotonic convergence for conditional expectations). *Let $\mathcal{G} \subseteq \mathcal{F}$ and $X_1, X_2, \dots \in \mathcal{L}^1$. Let one of the conditions apply:*

1. *Let $X \in \mathcal{L}^1$ such that $X_n \uparrow X$ is almost certain.*
2. *If $Y \in \mathcal{L}^1$ such that $|X_n| \leq |Y|$ for all n , and $X_n \xrightarrow{n \rightarrow \infty} X$ almost surely.*

Then

$$\mathbf{E}[X_n|\mathcal{G}] \xrightarrow{n \rightarrow \infty} \mathbf{E}[X|\mathcal{G}]$$

almost surely and in \mathcal{L}^1 .

Proof. For the \mathcal{L}^1 -convergence one has in both cases with theorem 11.2.3

$$\begin{aligned} \mathbf{E}[|\mathbf{E}[X_n|\mathcal{G}] - \mathbf{E}[X|\mathcal{G}]|] &= \mathbf{E}[|\mathbf{E}[X_n - X|\mathcal{G}]|] \\ &\leq \mathbf{E}[|X_n - X|] \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

We divide the almost certain convergence into the two cases: in case case 1. it is clear from Theorem 11.2.2 that $\mathbf{E}[X_n|\mathcal{G}]$ grows monotonically. Furthermore, for $A \in \mathcal{F}$ with the theorem of monotonic convergence

$$\mathbf{E}\left[\sup_n \mathbf{E}[X_n|\mathcal{G}]; A\right] = \sup_n \mathbf{E}[\mathbf{E}[X_n|\mathcal{G}]; A] = \sup_n \mathbf{E}[X_n; A] = \mathbf{E}[\sup_n X_n; A] = \mathbf{E}[X; A].$$

However, this shows that almost certainly $\sup_n \mathbf{E}[X_n|\mathcal{G}] = \mathbf{E}[X|\mathcal{G}]$ applies.

In case 2. we set

$$\begin{aligned} Y_n &:= \sup_{k \geq n} X_k \downarrow \limsup_n X_n = X \text{ almost certainly,} \\ Z_n &:= \inf_{k \geq n} X_k \uparrow \liminf_n X_n = X \text{ almost sure.} \end{aligned}$$

Thus $-Y \leq Z_n \leq X_n \leq Y_n \leq Y$, i.e. in particular $Y_1, Z_1, Y_2, Z_2, \dots \in \mathcal{L}^1$, so after 1.

$$\mathbf{E}[X|\mathcal{G}] = \lim_{n \rightarrow \infty} \mathbf{E}[Z_n|\mathcal{G}] \leq \lim_{n \rightarrow \infty} \mathbf{E}[X_n|\mathcal{G}] \leq \lim_{n \rightarrow \infty} \mathbf{E}[Y_n|\mathcal{G}] = \mathbf{E}[X|\mathcal{G}]$$

is almost certain. In particular, $\mathbf{E}[X_n|\mathcal{G}] \xrightarrow{n \rightarrow \infty} \mathbf{E}[X|\mathcal{G}]$ is almost certain. \square

11.3 The case $\mathcal{G} = \sigma(X)$

In the case $\mathcal{G} = \sigma(X)$, $\mathbf{E}[Y|X] := \mathbf{E}[Y|\sigma(X)]$ is the expectation of Y , given that the random variable random variable X is fixed. This is a function of X , as Proposition 11.7 shows.

Proposition 11.7 (condition on a random variable). *Let (Ω', \mathcal{F}') be a measurement space, X a Ω' -valued random variable with values in Ω' and $Y \in \mathcal{L}^1$. Then there exists a $\mathcal{F}'/\mathcal{B}(\mathbb{R})$ -measurable mapping $\varphi : \Omega' \rightarrow \mathbb{R}$ with $\mathbf{E}[Y|X] = \varphi(X)$.*

Proof. Clearly according to lemma 6.2 \square

Example 11.8 (Random probability of success). *Let us consider the question posed in Example 11.1 regarding the existence of the conditional probability $\mathbf{P}(Y = k|X)$, where X is uniform on $[0, 1]$ and X is independently binomially distributed with n and X . We now show that*

$$\mathbf{P}(Y = k|X) = \binom{n}{k} X^k (1 - X)^{n-k}. \quad (11.3)$$

Let $A = \{X \in I\}$ for $I \in \mathcal{B}([0, 1])$, i.e. A is a $\sigma(X)$ -measurable quantity. Then applies

$$\mathbf{E}[1_{Y=k}; A] = \mathbf{P}(Y = k, X \in I) = \int_I \binom{n}{k} x^k (1 - x)^{n-k} dx = \mathbf{E}\left[\binom{n}{k} X^k (1 - X)^{n-k}; A\right]$$

However, this means that (11.3) is true.

Example 11.9 (Sums of independent identically distributed random variables). *Let X_1, X_2, \dots be a sequence of independent, identically distributed random variables, $\mu = \mathbf{E}[X_1]$ and $S_n := X_1 + \dots + X_n$. Then*

$$\begin{aligned} \mathbf{E}[S_n|X_1] &= \mathbf{E}[X_1|X_1] + \mathbf{E}[X_2 + \dots + X_n|X_1] = X_1 + (n - 1)\mu, \\ \mathbf{E}[X_1|S_n] &= \frac{1}{n} \sum_{i=1}^n \mathbf{E}[X_i|S_n] = \frac{1}{n} \mathbf{E}[S_n|S_n] = \frac{1}{n} S_n. \end{aligned}$$

In the second calculation, for example, for $X = S_n$ and $Y = X_1$ the function φ from Proposition 11.7 is given by $\varphi(x) = \frac{1}{n}x$.

Example 11.10 (Buffon's needle problem). *On a plane there are straight lines at a horizontal distance of 1. needles of length 1 are thrown onto the plane; see Figure 2. Let us consider a needle. We set*

$$Z := \begin{cases} 1, & \text{if the needle intersects a straight line} \\ 0, & \text{otherwise} \end{cases}.$$

The center of the needle X is away from the left straight line and the extension of the needle




Figure 2: Sketch of Buffon's needle problem

makes an acute angle Θ with the left straight line. This means that X is uniform on $[0; 1]$, Θ is uniformly independent on $[0; \frac{\pi}{2}]$ and the following applies

$$\mathbf{P}(Z = 1|\Theta) = \mathbf{P}(X \leq \frac{1}{2} \sin(\Theta) \text{ or } X \geq 1 - \frac{1}{2} \sin(\Theta)) = \sin(\Theta).$$

This means that

$$\mathbf{P}(Z = 1) = \mathbf{E}[\mathbf{P}(Z = 1|\Theta)] = \mathbf{E}[\sin(\Theta)] = \frac{2}{\pi} \int_0^{\pi/2} (\sin(\theta)d\theta) = \frac{2}{\pi}.$$

This can be interpreted as follows: if you want to determine by simulation (i.e. by a Monte Carlo method) to find the numerical value of π you can simulate Buffon's needles. Since each individual needle has the probability $\frac{2}{\pi}$ of hitting a vertical line, is approximately

$$\pi \approx \frac{2}{\text{proportion of needles that hit a vertical line}}$$

according to the law of large numbers.

Example 11.11 (Search in lists). Given n names of people who come from r different cities. Each person comes (independently of any other) with probability p_j from city j , $j = 1, \dots, r$. The names (together with other personal data) are entered in r different (unordered) lists. If you now want a (random, according to (randomly distributed according to the probabilities p_1, \dots, p_r) in the list, first determine the list, you first determine the city j from which the person comes from and then search the list j for the person's name. Until you realize that the name does not appear in the list you have to compare the person to be found with names on the list to be found. The question now is: How many times on average have you the name of the person to be found with names on the list without success until you finally know that the person is not on the list?

We first define a few random variables:

J : number of the city from which the person to be searched comes

L : number of unsuccessful comparisons until the name of the person to be found is found

Z_j : number of people from city j

and $Z = (Z_1, \dots, Z_r)$. In order to determine $\mathbf{E}[L]$, we first determine we first determine

$$\mathbf{P}(L = a|J, Z) = 1_{Z_J=a}$$

and thus

$$\mathbf{P}(L = a|Z) = \sum_{j=1}^r p_j 1_{Z_j=a}.$$

From this we conclude

$$\mathbf{E}[L|Z] = \sum_{a=1}^{\infty} \sum_{j=1}^r a \cdot p_j \cdot 1_{Z_j=a} = \sum_{j=1}^r p_j Z_j$$

and therefore

$$\mathbf{E}[L] = \mathbf{E}[\mathbf{E}[L|Z]] = \sum_{j=1}^r p_j \mathbf{E}[Z_j] = n \cdot \sum_{j=1}^r p_j^2.$$

Example 11.12 (mixture of Poisson distributions). Let $\lambda > 0$ and $\lambda \sim \exp(\lambda)$ and for a given λ let $X \sim \text{Poi}(\lambda)$. We now show that $X + 1 \sim \text{geo}(1/(1 + \lambda))$.

Because: According to Proposition 9.25, the distribution is clearly determined by the characteristic function. First of all, the characteristic function of $Y \sim \text{geo}(p)$

$$t \mapsto \mathbf{E}[e^{itY}] = \sum_{k=0}^{\infty} (1-p)^{k-1} p e^{itk} = p e^{it} \sum_{k=0}^{\infty} ((1-p)e^{it})^k = \frac{p e^{it}}{1 - (1-p)e^{it}} = \frac{\frac{p}{1-p} e^{it}}{\frac{1}{1-p} - e^{it}}.$$

We calculate with example 6.13.2 for $t \in \mathbb{R}$

$$\mathbf{E}[e^{it(X+1)}] = e^{it} \mathbf{E}[\mathbf{E}[e^{itX} | \Lambda]] = e^{it} \mathbf{E}[e^{-(1-e^{it})\Lambda}] = \frac{\lambda e^{it}}{1 + \lambda - e^{it}},$$

so that the assertion with $\lambda = p/(1-p)$ or $p = 1/(1 + \lambda)$ follows.

Similarly, conditional independence is defined for

Markov chains; see for example at most countable and $P = (p_{ij})_{i,j \in E}$ a

We start in Proposition ?? with Doob's

According to proposition ??, $(|X_t|^p)_{t \in I}$ is

The first inequality is clear. For the second

1. it suffices to show the assertion for small $|x|$, since it is true for to show

We now come to 3. by showing $\mathbf{E}[T_n] = 1$. Because